

Serum concentrations of haptoglobin and serum amyloid A in water buffaloes (*Bubalus bubalis*) with abomasal ulcer

Javad Tajik^{1*}, Saeed Nazifi², Mahdi Heidari², Marzieh Babazadeh²

¹Department of Clinical Studies, School of Veterinary Medicine, Shahid Bahonar University of Kerman, Kerman, Iran; ²Department of Clinical Studies, School of Veterinary Medicine, Shiraz University, Shiraz, Iran

Article Info	Abstract
Article history: Received: 26 November 2011 Accepted: 20 February 2012 Available online: 15 September 2012	To evaluate the serum concentrations of haptoglobin (Hp) and serum amyloid A (SAA) in water buffaloes with abomasal ulcers, the abomasums of 100 randomly selected water buffaloes were examined after slaughter. Type I abomasal ulcers were found in 56 out of 100 buffaloes. Serum concentrations of Hp and SAA were measured. There was no significant difference between affected and non-affected buffaloes in the serum concentrations of Hp and SAA. The serum concentrations of Hp and SAA had no significant correlation with age and the serum SAA revealed no significant correlation with the number of abomasal ulcers. A significant correlation was found between the serum Hp and the number of abomasal ulcers ($r = 0.29, p = 0.04$). There was no significant difference in the serum concentrations of Hp and SAA between buffaloes with different ulcer locations in the abomasums. Although more work on a larger number of animals is required in this area, it seems that the measurement of the serum Hp can be used to predict the abundance of type I abomasal ulcers.
Key words: Haptoglobin Serum amyloid A Abomasal ulcer <i>Bubalus bubalis</i>	

© 2012 Urmia University. All rights reserved.

غلظت سرمی هاپتوگلوبولین و سرم آمیلوئید A در گاومیشهای مبتلا به زخم شیردان

چکیده

به منظور ارزیابی غلظت سرمی هاپتوگلوبولین و سرم آمیلوئید A در گاومیشهای مبتلا به زخم شیردان، پس از کشتار شیردان ۱۰۰ گاومیش که بصورت تصادفی انتخاب شده بودند بررسی گردید. زخم شیردان نوع یک در ۵۶ گاومیش مشاهده گردید. غلظت سرمی هاپتوگلوبولین و سرم آمیلوئید A اندازه گیری گردید. غلظت سرمی هاپتوگلوبولین و سرم آمیلوئید A بین گاومیشهای مبتلا و غیر مبتلا اختلاف معنی داری نداشت. همبستگی معنی داری بین غلظت سرمی هاپتوگلوبولین و سرم آمیلوئید A با سن دامها وجود نداشت و غلظت سرمی سرم آمیلوئید A همبستگی معنی داری با تعداد زخمهای شیردان نداشت. یک همبستگی معنی دار بین غلظت سرمی هاپتوگلوبولین و تعداد زخمهای شیردان مشاهده گردید ($r = 0.29, p = 0.04$). بین گاومیشهای با محل‌های مختلف زخم در شیردان اختلاف معنی داری از نظر غلظت سرمی هاپتوگلوبولین و سرم آمیلوئید A وجود نداشت. اگرچه مطالعات بیشتر که شامل دامهای بیشتری باشد مورد نیاز است، اما به نظر می رسد بتوان از غلظت سرمی هاپتوگلوبولین برای تشخیص تعداد زخم شیردان نوع یک در گاومیشهای مبتلا استفاده نمود.

واژه های کلیدی: هاپتوگلوبولین، سرم آمیلوئید A، زخم شیردان، گاومیش

*Correspondence:

Javad Tajik, DVM, DVSc
Department of Clinical Studies, School of Veterinary Medicine, Shahid Bahonar University of Kerman, Kerman, Iran.
E-mail: tajik@uk.ac.ir

Introduction

There are numerous reports regarding the occurrence of gastric ulcers in human, cattle, and swine. Different age groups of ruminants may be affected by abomasal ulcers.¹ Abomasal ulcers were divided into four types, each type producing distinct clinical signs. However, Type I ulcers, non-perforating erosions of abomasal mucosa, have mild or undetectable clinical signs and are usually not diagnosed until slaughter.² Braun *et al.* reported that 20.5 % of slaughtered cows had type I abomasal ulcer.¹ The results of Aukema and Breukink's study showed that 6.3 % of slaughtered cows had abomasal ulcers or ulcer scars.³ Although the etiology and pathogenesis of abomasal ulcers are not completely known, the clinical status of animals, nutrition, concurrent diseases and stress factors have been proposed as the probable factors causing the different prevalence rates of abomasal ulcers reported in different studies.^{1,4,5} Braun *et al.* believe that clinically normal animals have mainly, type I ulcers, whereas animals with clinical signs of disease often have type 2 to 4 ulcers.¹ Although the economic effect of abomasal erosions in ruminants has not yet been investigated, physical discomfort resulting in reduced feed consumption and weight gain have been proposed as the probable causes of loss.⁶

Iranian water buffalo (*Bubalus bubalis*) has a high economic value by providing meat, milk and labor for local farmers. Ghadrnan-Mashhadi *et al.* reported a high prevalence of type I abomasal ulcers (63.5 %) in slaughtered water buffaloes in Iran.⁷ Despite the probable importance of abomasal ulcers in water buffalo in Iran, there is a little information about the clinical signs and no previous study regarding the diagnosis of the abomasal ulcers in this species has been recorded.

Acute phase proteins (APPs) are a group of blood proteins that show changes in their concentration in animals subjected to external or internal challenges such as infection, inflammation, surgical trauma, or stress.⁸ It has been suggested that APPs may provide valuable diagnostic information in detection, prognosis and monitoring the diseases in animal species and animal health.⁹

Despite the growing interest in using APPs as health indicators,¹⁰ to the best of our knowledge, there has been no previous study regarding the probable changes of the APPs in abomasal ulcers in water buffalo. Therefore, the aim of this study was to evaluate and to compare the rate of serum haptoglobin (Hp) and serum amyloid A (SAA) between affected and non-affected water buffalo.

Materials and Methods

Animals and sampling. The investigation was carried out on water buffaloes (*Bubalus bubalis*) which were slaughtered in a slaughter house reserved only for buffaloes in Ahvaz City, southwestern Iran, from December 2009 to January 2010, and April to May 2010.

After clinical examination and before slaughter, jugular blood samples were collected from 100 clinically healthy water buffaloes in plane tubes, without anticoagulant. Immediately after slaughter, the abomasum of the animals were opened along the greater curvature and washed in water to free ingesta. The abomasal mucosa was examined, and the type, number, and location of the lesions were recorded. The most frequent type of lesion was referred to as the main lesion.¹ Buffaloes were of both sexes with different ages, and were selected randomly. The age of the animals was estimated using dental characteristics. All animals had grazed the previous summer on ranges around the city. The blood serum was separated after centrifugation at 1800 *g* for 10 min and the serum samples stored at -20 °C until analysis.

Assays. Total serum protein (TP) was measured by the Biuret method and serum concentration of Hp was measured according to prevention of the peroxidase activity of hemoglobin, which is directly proportional to the amount of Hp.¹⁰ The analytical sensitivity of this test in serum has been determined as 0.01 mg mL⁻¹ for Hp by the manufacturer (Tridelta Development Plc, Wicklow, Ireland).

Serum SAA was measured by a solid phase sandwich ELISA.¹⁰ The analytical sensitivity of this test in serum has been determined as 0.30 µg mL⁻¹ for SAA by the manufacturer (Tridelta Development Plc, Wicklow, Ireland).

Statistical analysis. Statistical analysis was performed using SPSS version 12 (Illinois, Chicago). The correlation of the measured serum parameters with age were analyzed by Pearson's correlation tests. Two sample *t*-tests were used to detect differences in the parameters between the two sexes and between affected and non-affected buffaloes. The buffaloes were divided into three groups, according to their age as G₁ ≤ 2 years, 2 years < G₂ ≤ 5 years, and G₃ > 5 years. Analysis of variance (ANOVA) tests were used to compare the serum proteins between the different age groups of water buffaloes. Differences were considered significant at *p* < 0.05.

Results

Overall, 69 male buffaloes and 31 female buffaloes were sampled. The average ages (Mean ± SEM) of the male and female buffaloes were 2.19 ± 0.10 and 2.68 ± 0.30 years, respectively. There was no significant difference between the two sexes in age.

The results of the serum concentration of measured proteins in affected and non-affected buffaloes and in both sexes are shown in Table 1.

Fifty buffaloes were examined in each sampling period and 43 and 20 buffaloes were diagnosed as affected in the first and second sampling periods, respectively.

All abomasal lesions were classified as type I abomasal ulcers, and were confirmed in histopathological examinations.

There was no significant difference between affected and non-affected buffaloes in the serum concentrations of Hp and SAA ($p > 0.05$). There was also no significant difference between the affected and non-affected buffaloes in age (2.45 ± 0.16 and 2.07 ± 0.16 , respectively) ($p > 0.05$). The serum concentrations of Hp and SAA had no significant correlation with age and the serum SAA revealed no significant correlation with the number of abomasal ulcers, however, serum Hp showed a significant correlation with the number of abomasal ulcers ($r=0.29$, $p=0.04$).

There was no significant difference in the serum concentrations of Hp and SAA between buffaloes with different ulcer locations in the abomasums and between different age groups ($p > 0.05$).

Table 1. The concentrations (Mean \pm SEM) of serum Hp and SAA in affected and non-affected buffaloes and in different genders and age groups.

Groups	No.	TP (g dL ⁻¹)	Hp (g L ⁻¹)	SAA (μ g mL ⁻¹)
All sampled	100	7.16 \pm 0.06	0.57 \pm 0.06	4.11 \pm 0.38
Affected	63	7.08 \pm 0.07	0.56 \pm 0.06	4.27 \pm 0.44
Non-affected	37	7.32 \pm 0.10	0.62 \pm 0.19	3.18 \pm 0.45
Male	69	7.15 \pm 0.07	0.55 \pm 0.07	4.17 \pm 0.54
Female	31	7.18 \pm 0.12	0.61 \pm 0.09	4.03 \pm 0.49
G ₁	64	7.21 \pm 0.08	0.53 \pm 0.07	4.16 \pm 0.46
G ₂	33	7.09 \pm 0.10	0.71 \pm 0.14	4.27 \pm 0.81
G ₃	3	6.98 \pm 0.29	0.35 \pm 0.12	2.84 \pm 0.79

Both sexes were evaluated separately, and the results of the comparison of the measured serum factors between the affected and non-affected buffaloes showed no difference ($p > 0.05$).

Different age groups of buffaloes were also evaluated separately and the results of the comparison of the measured serum factors between the affected and non-affected buffaloes showed no difference ($p > 0.05$).

Discussion

The reference value of serum Hp and SAA in healthy water buffaloes has been investigated and the results of Ghadr-dan-Mashhadi *et al.* and the current studies have revealed the high prevalence of type I abomasal ulcers in water buffaloes in Iran.^{7,11} However, to the best of our knowledge, there has been no previous research regarding the changes of the serum Hp and SAA in abomasal ulcers in water buffaloes. The mean serum concentration of Hp for healthy water buffaloes in the current study (Mean \pm SEM: 0.62 ± 0.19) was slightly higher than that previously reported as the normal ranges for healthy water buffaloes ($0.13 - 0.06$ g L⁻¹) by Tajik *et al.*¹¹ According to our results, the mean serum concentration of SAA for healthy water buffaloes was 3.18 ± 0.45 μ g mL⁻¹ and was slightly higher than the previously reported mean serum concentration

of SAA in healthy water buffaloes (2.99 ± 0.14 μ g mL⁻¹).¹¹ Some factors such as age, sex, breed, season, geographic and dietary factors affect the serum proteins,^{12,13} which may cause the observed differences.

Following tissue destruction and inflammation, change in blood serum proteins is expected.¹⁴ During inflammation, the protein production in liver is switched towards increased synthesis of positive acute phase proteins such as Hp and SAA.^{8,14,15} Haptoglobin is synthesized in the liver and is a major acute-phase protein in numerous species of animals.^{16,17} It is believed that the circulating level of Hp in ruminants is negligible in normal animals, however, increases over 100-fold with immune stimulation.^{18,19} Serum amyloid A is also considered as one of the major acute phase reactants in vertebrate,¹⁰ and it is generally assumed that SAA have a protective role during inflammation.²⁰ According to our results, although there was no significant difference between affected and non-affected buffaloes in the serum concentrations of Hp and SAA, there was a significant correlation between the serum Hp and the number of abomasal ulcers. All detected abomasal ulcers in examined buffaloes were classified as type I, non-perforating erosions of abomasal mucosa, which are mucosal defects and do not penetrate the deeper layers of abomasum.⁵ It seems that little penetration of type I abomasal ulcer to the abomasal layers with no consequences, such as peritonitis (which occurs in type III and IV abomasal ulcer) or bleeding (which occurs in type II abomasal ulcer), causes a negligible effect on the blood serum Hp and SAA concentrations. However, the observed positive correlation between the number of abomasal ulcers and the serum Hp may be due to the moderate inflammatory reactions at the ulcer places, which were confirmed in the histopathological examination.

We found no significant difference between either sex in the serum concentrations of the measured acute-phase proteins, which was the same as that reported previously in water buffalo and other species, such as cattle and sheep.^{11,21,22}

Our results also showed that serum Hp and SAA had no significant correlation with age and there was no significant difference between different age groups. Similar results have been found in previous studies in water buffalo, sheep and cattle.^{11,21,22} However, age-related changes in serum concentrations of Hp and SAA have been reported in horse.²³⁻²⁵ It is believed that species variations in APPs occur due to inherent variations in synthesis and in rates of entry and exit of these APPs from circulation.²⁶

It has been shown that the occurrence and severity of inflammatory diseases in different species of domestic animals can be evaluated by acute phase protein profiling.^{10,17,27} It is believed that the stage of disease can be better evaluated by monitoring more than one acute phase protein.¹⁵ The findings of the present study showed that although measurement of serum Hp and SAA cannot

be used solely in the diagnosis of type I abomasal ulcers in water buffaloes, measurement of the serum Hp can be used in prediction of the abundance of the type I abomasal ulcers. Different affected abomasal layers and distinctive consequences of the different types of abomasal ulcers such as local peritonitis in type III abomasal ulcers cause their effects on the serum proteins to be different and incomparable. It seems that more work on a larger number of animals and in other types of abomasal ulcers is needed before these findings can be used in practice.

References

- Braun U, Eicher R, Ehrensperger F. Type 1 Abomasal Ulcers in Dairy Cattle. *J Vet Med A* 1991; 38: 357-366.
- Whitlock RH. Bovine stomach diseases. In: Anderson NV, ed. *Veterinary Gastroenterology*. Philadelphia: Lea and Febiger 1980; 413-428.
- Aukema JJ, Breukink HJ. Abomasal ulcer in adult cattle with fatal hemorrhage. *Cornell Vet* 1974; 64: 303-317.
- Dirksen GU. Ulceration, dilation and incarceration the abomasum in calves: Clinical investigations and experiences. *Bovine Pract* 1994; 28: 127-135.
- Smith DF, Munson L, Erb HN. Abomasal ulcer disease in adult dairy cattle. *Cornell Vet* 1983; 73: 213-224.
- Jensen R, Spraker TR, Glock RD, et al. Abomasal erosions in feedlot cattle. *Am J Vet Res* 1992; 53:110-115.
- Ghadrdan-Mashhadi A, Esmaeilzadeh S, Gandomi M. Prevalence of different abomasal ulcers in buffalos of Ahvaz: an abatorial survey. *J Vet Res* 2010; 65: 351-356.
- Murata H, Shimada N, Yoshioka M. Current research on acute phase proteins in veterinary diagnosis: an overview. *Vet J* 2004; 168: 24-40.
- González FH, Tecles F, Martínez-Subiela S, et al. Acute phase protein response in goats. *J Vet Diagn Invest* 2008; 20: 580-584.
- Baghshani H, Nazifi S, Saeb M, et al. Influence of road transportation on plasma concentrations of acute phase proteins, including fibrinogen, haptoglobin, serum amyloid A, and ceruloplasmin, in dromedary camels (*Camelus dromedarius*). *Comp Clin Pathol* 2010; 19: 193-198.
- Tajik J, Nazifi S, Eskandarzadeh N, et al. Serum haptoglobin, amyloid A, and sialic acid values in healthy water buffalo. *Online J Vet Res* 2011; 15: 270-276.
- Khadjeh GH, Razi Jalali M. Electrophoretic pattern of blood serum proteins in the Khuzestan indigenous buffaloes. *J Fac Vet Med Univ Tehran* 2003; 58: 307-311.
- Mostaghni K, Nazifi S, Soleymani D. A study of CSF and serum proteins in sheep by electrophoresis. *J Fac Vet Med Univ Tehran* 1996; 50: 69-77.
- Saleh MA, Rateb HZ, Misk NA. Comparison of blood serum proteins in water buffaloes with traumatic reticuloperitonitis and sequelae. *Res Vet Sci* 2008; 85: 208-213.
- Eckersall PD. Proteins, Proteomics, and the Dysproteinemias. In: Kaneko JJ, Harvey JW, Bruss ML, Eds. *Clinical Biochemistry of Domestic Animals*. 6th ed. New York: Academic Press Inc. 2008; 117-155.
- Feldman BF, Zinkl JG, Jain NC. *Schalm's Veterinary Hematology*. Philadelphia: Lippincott Williams & Wilkins 2000; 891-896.
- Nazifi S, Saeb M, Ghasemian O, et al. Evaluation of serum haptoglobin in clinically healthy Iranian camels (*Camelus dromedarius*). *Comp Clin Pathol* 2006; 15:195-197.
- Conner JG, Eckersall PD, Wiseman A, et al. Bovine acute phase response following turpentine injection. *Res Vet Sci* 1988; 44: 82-88.
- Conner JG, Eckersall PD, Wiseman A, et al. Acute phase response in calves following infection with *Pasteurella haemolytica*, *Ostertagia ostertagi* and endotoxin administration. *Res Vet Sci* 1989; 47:203-207.
- Uhlar CM, Whitehead AS. Serum amyloid A, the major vertebrate acute-phase reactant. *Eur J Biochem* 1999; 265:501-523.
- Nazifi S, Rezakhani A, Koohimoghadam M, et al. Evaluation of serum haptoglobin in clinically healthy cattle and cattle with inflammatory diseases in Shiraz, a tropical area in southern Iran. *Bulg J Vet Med* 2008; 11: 95-101.
- Nowroozi-Asl A, Nazifi S, Bahari A. Determination of serum haptoglobin reference value in clinically healthy Iranian fat-tailed sheep. *Iran J Vet Res* 2008; 9:171-173.
- Harvey JW, Asquith RL, McNulty PK, et al. Hematology of foals up to one year old. *Equine Vet J* 1984; 16: 347-353.
- Paltrinieri S, Giordano A, Villani M, et al. Influence of age and foaling on plasma protein electrophoresis and serum amyloid A and their possible role as markers of equine neonatal septicemia. *Vet J* 2008; 176: 393-396.
- Taira I, Fujinaga T, Okumura M. Equine haptoglobin isolation, characterization, and the effects of ageing, delivery, inflammation on the serum concentration. *J Vet Med Sci* 1992; 54: 435-442.
- Kaneko JJ. Serum proteins and dysproteinemias. In: Kaneko JJ, Harvey JW, Bruss M, Eds. *Clinical biochemistry of domestic animals*, 5th ed. San Diego: Academic Press Inc 1997; 117-138.
- Horadagoda A, Edwards SER, Eckersall PD. Acute phase proteins in cattle: discrimination between acute and chronic inflammation. *Vet Rec* 1999; 144: 437-441.