

Effect of extracellular matrix on testosterone production during *in vitro* culture of bovine testicular cells

Vahid Akbarinejad¹, Parviz Tajik^{2*}, Mansoureh Movahedin³, Reza Youssefi²

¹ Young Researchers and Elites Club, Roudehen Branch, Islamic Azad University, Roudehen, Iran; ² Department of Theriogenology, Faculty of Veterinary Medicine, University of Tehran, Tehran, Iran; ³ Department of Anatomical Sciences, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran.

Article Info

Article history:

Received: 28 September 2015
Accepted: 05 March 2016
Available online: 15 March 2017

Key words:

Bovine testicular cells
Extracellular matrix
Testosterone

Abstract

Testosterone is believed to play a significant role in spermatogenesis, but its contribution to the process of spermatogenesis is not completely understood. Given that extracellular matrix (ECM) facilitates differentiation of spermatogonial stem cells (SSCs) during culture, the present study was conducted to elucidate whether testosterone contribute to the permissive effect of ECM on SSCs differentiation. In experiment 1, testosterone production was measured in testicular cells cultured for 12 days on ECM or plastic (control). In experiment 2, testosterone production was assessed in testicular cells cultured on ECM or plastic (control) and exposed to different concentrations of hCG. In experiment 3, the gene expression of factors involved in testosterone production was analyzed. Testosterone concentration was lower in ECM than in the control group in experiment 1 ($p < 0.05$). In experiment 2, testosterone concentration was increased in response to hCG in both groups but cells cultured on ECM were more responsive to hCG than those cultured on plastic ($p < 0.05$). In the experiment 3, qRT-PCR revealed the inhibitory effect of ECM on the gene expression of steroidogenic acute regulatory protein (StAR) ($p < 0.05$). Nevertheless, the expression of LH receptor was greater in ECM-exposed than in unexposed cells ($p < 0.05$). In conclusion, the present study showed that inhibiting the expression of StAR, ECM could lower testosterone production by Leydig cells during *in vitro* culture. In addition, the results indicated that ECM could augment the responsiveness of Leydig cells to hCG through stimulating the expression of LH receptor.

© 2017 Urmia University. All rights reserved.

تأثیر ماتریکس خارج سلولی بر تولید تستوسترون در کشت برون زیستی سلول‌های بیضه گاو

چکیده

تستوسترون نقش کلیدی در اسپرماتوژنز ایفا می‌کند، ولی اطلاعات کاملی در این رابطه در دست نیست. نظر به اینکه ماتریکس خارج سلولی (ECM) تمایز سلول‌های بنیادی اسپرماتوگونیال (SSCs) را در طول کشت تسهیل می‌کند، مطالعه حاضر جهت مشخص کردن نقش تستوسترون در رابطه با تأثیر ECM بر تمایز SSCs به انجام رسید. در آزمایش ۱، تولید تستوسترون در سلول‌های بیضه کشت داده شده بر روی ECM و پلاستیک (گروه شاهد) به مدت ۱۲ روز مورد ارزیابی قرار گرفت. در آزمایش ۲، میزان تولید تستوسترون در سلول‌های بیضه که بر روی ECM و پلاستیک کشت داده شده بودند و با مقادیر مختلف هورمون کوریونی انسانی (hCG) مواجه شده بودند، اندازه‌گیری شد. در آزمایش ۳، بیان ژن فاکتورهای دخیل در تولید تستوسترون ارزیابی شد. تولید تستوسترون در آزمایش ۱ در گروه ECM کمتر از گروه شاهد بود ($p < 0.05$). در آزمایش ۲، تولید تستوسترون در پاسخ به hCG در هر دو گروه افزایش پیدا کرد، اما سلول‌هایی که بر روی ECM کشت داده شده بودند در مقایسه با سلول‌هایی که بر روی پلاستیک کشت داده شده بودند نسبت به hCG حساس‌تر بودند ($p < 0.05$). در آزمایش ۳، qRT-PCR نشان دهنده نقش مهار ECM بر بیان ژن پروتئین تنظیم کننده استروئیدوژنز (StAR) بود ($p < 0.05$). با این وجود، بیان ژن گیرنده LH در گروه ECM در مقایسه با گروه شاهد بالاتر بود ($p < 0.05$). در نتیجه، مطالعه حاضر نشان داد که ECM با مهار بیان StAR می‌تواند تولید تستوسترون توسط سلول‌های لیدیگ را در طول کشت برون زیستی کاهش دهد. به علاوه، نتایج مشخص کرد که ECM می‌تواند با تحریک بیان ژن گیرنده LH حساسیت سلول‌های لیدیگ را به hCG افزایش دهد.

واژه‌های کلیدی: تستوسترون، سلول‌های بیضه گاو، ماتریکس خارج سلولی

*Correspondence:

Parviz Tajik. DVM, DVSc
Department of Theriogenology, Faculty of Veterinary Medicine, University of Tehran, Tehran, Iran.
E-mail: ptajik@ut.ac.ir

Introduction

Spermatogenesis is a complex process of cell proliferation and differentiation culminating in production of mature spermatozoa from spermatogonial stem cells (SSCs).^{1,2} Somatic cells encompassing SSCs, called niche cells, regulate the self-renewal and differentiation of SSCs by production of hormones and growth factors, thereby contributing to the perpetuation of spermatogenesis.¹

Testosterone, produced by Leydig cells in the testis, is one of the factors that is believed to play an essential role in spermatogenesis^{3,4} as some men with diminished spermatogenesis experience low levels of circulatory testosterone concentration as well.⁵ Dysfunction of Leydig cells has also been observed to lead to spermatogenic disruption.^{6,7} In addition, induction of spermatogenesis in mice with gonadotropin deficiency has been reported following treatment with testosterone.⁸ Indeed, three major defects have been determined to ensue testosterone deficiency and/or disruption in testosterone action including disintegration of blood-testis barrier,⁹⁻¹² blockade in conversion of round to elongated spermatids¹³⁻¹⁵ and disruption in mature spermatozoa release from Sertoli cells.¹³ Yet the role of testosterone in the process of spermatogenesis is not completely understood and requires to be elucidated.⁴

Testosterone synthesis by Leydig cells is under the influence of luteinizing hormone (LH). This mechanism could also be regulated by the LH analogue, human chorionic gonadotropin (hCG).¹⁶ Steroidogenic acute regulatory protein (StAR), which is produced in response to LH, plays an essential role in testosterone production through translocating intracellular cholesterol to inner mitochondrial membrane in which the cholesterol side chain would be cleaved leading to production of pregnenolone, the precursor for testosterone.¹⁷

Extracellular matrix (ECM), composed of macromolecules with different structural and functional properties, could impact the SSCs function through generation of signals by ECM molecules *per se*, as well as modulating the other extracellular stimuli, including growth and differentiating factors.¹⁸⁻²⁰ Application of ECM during *in vitro* culture of cells provides an environment partly resembling the normal tissue^{18,21,22} and has been indicated to have permissive effect for differentiation of SSCs.²³⁻²⁵ This phenomenon has been attributed to the impact of ECM on the behavior of niche cells and the interaction between niche and germ cells.²⁴ The ECM has been observed to influence the shape and function of Leydig cells.²⁶ In addition, ECM has been reported to influence the production of testosterone by Leydig cells.²⁷⁻²⁹

In the previous studies investigating the effect of ECM on SSCs differentiation, the impact of ECM on the potential contribution of testosterone in terms of SSCs differentiation has not been assessed.²³⁻²⁵ Therefore, the

present study was conducted to address whether testosterone contribute to the differentiating effect of ECM on SSCs during *in vitro* culture.

Materials and Methods

Animals and testicular biopsy. Animal Ethics Committee at University of Tehran approved the present study in terms of animal welfare and ethics. Testicular biopsies were obtained from Holstein calves (n = 17; aged 3 to 5 months) as previously described.²³ In brief, testicular biopsy was performed under sedation with xylazine (0.20 mg kg⁻¹; Alfasan, Woerden, Holland) and local anesthesia with lidocaine (Aburaihan Pharmaceutical Co., Tehran, Iran). Following incision, the testicular tissue was obtained and placed into a 15 mL tube containing Dulbecco' minimal essential medium (DMEM; Gibco, Carlsbad, USA) with 10% fetal bovine serum (FBS; Sigma-Aldrich, St. Louis, USA) and antibiotics (100 IU mL⁻¹ penicillin and 100 µg mL⁻¹ streptomycin; Gibco). Following testicular biopsy, the specimen was transferred on ice to the laboratory within 2 hr.

Experiment 1. Testicular cells (n = 7 cell populations from different calves) were cultured for 12 days and testosterone production was measured on days 6 and 12 of culture.²³

Experiment 2. Testicular cells (n = 7 cell populations from different calves) were cultured for 7 days. On day 6, hCG (IVF-C®; LG Life Sciences, Seoul, South Korea) was added to cultures at concentrations of 0.00, 0.10, 1.00 and 10.00 IU L⁻¹ for a one-day period, afterwards testosterone concentration was evaluated.²⁷

Experiment 3. Testicular cells (n = 3 cell populations from different calves) were cultured for 12 days and gene expression of luteinizing hormone receptor (LHR), CYP12A1, which codes for a member of the cytochrome P450 superfamily of enzymes contributing to androgen synthesis³⁰ and steroidogenic acute regulatory protein (StAR), which is responsible for transferring cholesterol into mitochondria thereby contributing to steroidogenesis,³¹ were evaluated using quantitative real time PCR on days 6 and 12.²³

Cell isolation. Cell isolation was implemented using a two-step enzymatic isolation procedure.²³ The testicular tissue was washed three times in DMEM containing antibiotics and was minced into small pieces using a sterile scissor. The minced testicular tissue was incubated in DMEM containing 1 mg mL⁻¹ collagenase (Sigma-Aldrich), 1 mg mL⁻¹ hyaluronidase (Sigma-Aldrich), 1 mg mL⁻¹ trypsin (Sigma-Aldrich) and 5 µg mL⁻¹ DNase (Fermentas, Waltham, USA) at 37 °C in a shaker incubator with 80 cycles per minute for approximately 60 min. The digested testicular tissue was washed three times with DMEM and the supernatant was disposed after each washing, leading to isolation of seminiferous tubules. During the second step

of enzymatic digestion, seminiferous tubules were incubated at 37 °C in DMEM containing 1 mg mL⁻¹ collagenase, 1 mg mL⁻¹ hyaluronidase and 5 µg mL⁻¹ DNase until disintegration of the seminiferous tubules and separation of the constituent cells. Individual cells were isolated from the remaining tubule fragments by centrifugation at 30 *g* for 2 min. Following filtration through 77 and 55 mm nylon filters, the cells were pelleted. The pellet was re-suspended in the DMEM containing antibiotics and 10% knock-out serum replacement (KSR; Gibco).

Cell culture. Wells used for the control group were uncoated. Wells used for the ECM group were coated with ECM gel (Sigma-Aldrich) as the manufacturer indicated. The ECM gel was prepared from Engelbreth-Holm-Swarm (EHS) mouse sarcoma and was composed of laminin as the major component, collagen type IV, heparan sulfate proteoglycan, entactin and other minor components. In the experiments related to assessment of gene expression and testosterone concentration, 6 and 24-well plates were used, respectively. In the experiments associated with evaluation of gene expression and testosterone measurement, cells were seeded at concentrations of 1 × 10⁶ and 3 × 10⁵ cells per well containing DMEM with antibiotics and 10% KSR, respectively. The plates were incubated at 37 °C in a humidified atmosphere with 5% CO₂. The medium was replaced with fresh one every three days.²³

Testosterone concentration analysis. Testosterone concentration was measured by direct, competitive, chemiluminescence immunoassay kit (DiaSorin, Stillwater, USA).

RNA isolation and quantitative real-time PCR (qRT-PCR). Following trypsinization of the cultured cells, the isolated cells were subjected to total RNA extraction using Trizol reagent (Fermentas). The extracted RNA was treated with DNase (Fermentas) to eliminate DNA contamination. The concentration of RNA was determined using UV spectrophotometry (Eppendorff, Hamburg, Germany). The cDNAs were synthesized from 500 ng of RNA by oligo (dT) primers using RevertAid™ first strand cDNA synthesis kit (Fermentas). Primers for genes of interest are shown in Table 1. The PCRs were performed using master mix and SYBR Green I (Fermentas) in a thermal cycler (Applied Biosystems, California, USA). The PCR program started with an initial melting cycle for 5 min at 95 °C to activate the polymerase, followed by 40 cycles of melting (30 sec at 95 °C), annealing (30 sec at 58 °C) and extension (30 sec at 72 °C). The quality of the PCR reactions

was confirmed by melting curve analyses. For each sample, the reference gene (*β-actin*) and target gene were amplified in the same run. The target genes were normalized to the reference gene. The mean target gene threshold cycle (Ct) and mean exogenous control (*β-actin*) Ct for each sample were calculated from duplicate wells. The target gene Ct of the control was subtracted from the Ct of target gene, resulting in ΔCt. In each experiment, the Ct of time-point 0 sample was considered as calibrator. Subsequently, the ΔCt of sample was then subtracted from the ΔCt of calibrator, resulting in the ΔΔCt, which was used for calculation of the relative amounts of target gene expression for each sample.³²

Statistical analysis. Initially, datasets were tested for normal distribution using Kolmogorov-Smirnov test (univariate procedure). Data associated with testosterone concentration in experiment 1 were log (base 10) transformed prior to analysis because of the lack of normal distribution. Data were analyzed using mixed procedure including random and repeated statements in the model to specify co-variation between and within calves respectively.³³ In addition, LSMEANS statement was used to perform multiple comparisons. All analyses were conducted in SAS (version 9.2; SAS Institute, Cary, USA). Data are presented as mean ± SD. Differences with *p* < 0.05 were considered significant.

Results

Experiment 1. Testosterone concentration was higher in the control than ECM group on days 6 and 12 of culture (*p* < 0.05). However, the production of testosterone was not different between days 6 and 12 in the control and ECM groups (*p* > 0.05; Table 2).

Experiment 2. In the control group, testosterone concentration in cells cultured with 10.00 ng mL⁻¹ hCG was greater than that in cells cultured with 0.00 and 0.10 IU mL⁻¹ of hCG (*p* < 0.001). However, the concentration of testosterone did not differ among 0.00, 0.10 and 1.00 IU mL⁻¹ concentrations of hCG and between 1.00 and 10.00 IU mL⁻¹ concentrations of hCG (*p* > 0.05; Table 3). In ECM group, testosterone concentration in 10.00 IU mL⁻¹ hCG group was greater than that in 0.00, 0.10 and 1.00 IU mL⁻¹ hCG groups (*p* < 0.05). In addition, testosterone concentration was higher in 1.00 IU mL⁻¹ hCG group than 0.00 and 0.10 IU mL⁻¹ hCG groups (*p* < 0.05). Testosterone concentration was not different between 0.00 and 0.10 IU mL⁻¹ hCG groups (*p* > 0.05; Table 3).

Table 1. Primer sequences used for qRT-PCR.

Gene	Forward primer (5'-3')	Reverse primer (5'-3')
<i>β-actin</i>	TCG CCC GAG TCC ACA CAG	ACC TCA ACC CGC TCC CAA G
<i>Lhr</i> ^a	TGACCATGGCCCGTCTAAAA	TACTACCCAAAGCAATTTATAGATTCAATG
<i>Cyp17a1</i>	CACCGATATTATCAGAAAACCC	ATTGGTGATGGACTCAAAGG
<i>Star</i> ^b	GTGGAATCCCAATGTCAAGG	TGATGACCGTGTCTTTTCCA

^aLH receptor; ^bSteroidogenic acute regulatory protein.

Table 2. Testosterone concentration (ng mL⁻¹) in control and extracellular matrix (ECM) groups on Days 6 and 12 of culture. Data are presented as mean ± SD.

Gene	Day 6	Day 12
Control	3.50 ± 1.18 ^a	3.72 ± 1.70 ^a
ECM	1.64 ± 0.48 ^b	1.94 ± 0.77 ^b

^{ab} Different superscripts indicate significant differences within each column ($p < 0.05$).

Experiment 3. In the control group, the gene expression of LHR on day 6 (2.47 ± 0.62 fold) was higher than that on day 0 ($p < 0.05$); however, LHR expression on day 12 was not different from days 0 and 6 ($p > 0.05$). In ECM group, the expression of LHR on days 6 (5.00 ± 1.23 fold) and 12 (3.81 ± 1.38 fold) was greater than that on day 0 ($p < 0.01$), but it did not differ between days 6 and 12 ($p > 0.05$). On day 6, the gene expression of LHR was higher in ECM than the control group ($p < 0.05$), but it was not different between two groups on day 12 of culture ($p > 0.05$; Fig. 1).

The gene expression of CYP12A1 did not change over the course of culture in the control and ECM groups ($p > 0.05$) and it was not different between two groups on days 6 and 12 ($p > 0.05$; Fig. 2).

In the control group, the expression of StAR did not change during culture ($p > 0.05$). In ECM group, however, the expression of StAR was 77.00% and 70.00% lower on days 6 and 12, respectively, as compared to day 0 ($p < 0.05$). The expression of StAR in ECM group was lower than that in the control group on days 6 and 12 ($p < 0.05$; Fig. 3).




Fig. 1. Relative gene expression of LHR in the control and ECM groups (n = 3) on days 6 and 12. Lowercase letters (a and b) indicate significant difference within groups between different timepoints ($p < 0.05$). Uppercase letters (A and B) indicate significant difference between two experimental groups at the specified timepoint ($p < 0.05$).

Table 3. Testosterone concentration (ng mL⁻¹) in control and extracellular matrix (ECM) groups following exposure by different levels of hCG. Data are presented as mean ± SD.

Groups	hCG concentration (IU L ⁻¹)			
	0.00	0.10	1.00	10.00
Control	3.64 ± 0.65 ^{aA}	3.73 ± 0.89 ^{aA}	4.10 ± 0.74 ^{abA}	4.49 ± 0.70 ^{bA}
ECM	2.43 ± 0.44 ^{aB}	2.62 ± 0.43 ^{aB}	3.05 ± 0.39 ^{bb}	3.47 ± 0.38 ^{cB}

^{abc} Different lowercase superscripts indicate significant differences within each row ($p < 0.05$).

^{AB} Different uppercase superscripts indicate significant differences within each column ($p < 0.05$).




Fig. 2. Relative gene expression of CYP17A1 in the control and ECM groups (n = 3) on days 6 and 12. No significant differences were noted among the groups.




Fig. 3. Relative gene expression of StAR in the control and ECM groups (n = 3) on days 6 and 12. Lowercase letters (a and b) indicate significant difference within groups between different time points ($p < 0.05$). Uppercase letters (A and B) indicate significant difference between two experimental groups at the specified time point ($p < 0.05$).

Discussion

The present study revealed that ECM decreased the production of testosterone by Leydig cells during *in vitro* culture. Culturing testicular germ cells of rat on ECM for 22 days, Lee *et al.* reported differentiation of germ cells up to haploid cells.²⁴ The same observation was indicated when ECM was applied for culture of human testicular germ cells.²⁵ More recently, culturing bovine SSCs on ECM for a 12-day period, Akbarinejad *et al.* found evidence for the initiation of SSCs differentiation. This phenomenon was not observed following culture of SSCs on plastic.²³ On the other hand,

giving mice flutamide, an androgen receptor antagonist, Shetty *et al.* observed stimulation of differentiation in spermatogonia in mice.³⁴ The same phenomenon was detected when GnRH antagonist, decreasing intratesticular testosterone concentration, was used.³⁴ Further, it was revealed that combination of flutamide and GnRH antagonist intensified spermatogonial differentiation, whereas combination of testosterone with GnRH antagonist inhibited spermatogonial differentiation.³⁴ Taken together, it was concluded that testosterone played an inhibitory role in differentiation of spermatogonia through binding to androgen receptor,³⁴ which was substantiated by a further study.³⁵ Moreover, using a Sertoli-cell androgen receptor knockout model, Zhou *et al.* demonstrated the inhibitory effect of androgen receptor of Sertoli cells in spermatogonial differentiation.³⁶ Therefore, inhibition of testosterone synthesis by Leydig cells might be one of the mechanisms whereby ECM molecules facilitate SSCs differentiation.²³⁻²⁵ In our previous study, we indicated that ECM molecules mediated their differentiative effect on SSCs through upregulation of KIT ligand.²³ Further studies are warranted to elucidate the other potential mechanisms contributing to permissive effect of ECM on SSCs differentiation.

Evaluating the effect of different types of proteins composing ECM, Diaz *et al.* indicated that laminin did not influence testosterone production, however, collagen type IV and fibronectin decreased the production of testosterone by Leydig cells.²⁷ Further, investigating the downstream mechanisms whereby collagen type IV reduces testosterone production, Diaz *et al.* found that the inhibitory effect of collagen type IV was mediated through activating extracellular signal-regulated kinase and decreasing cAMP production and StAR expression.³⁷ The gene expression of StAR has also been revealed to be lower in the ECM-exposed cells compared to unexposed ones in the present study. Collagen type IV was a part of the ECM used in the present study. Therefore, the lower concentration of testosterone in ECM-coated plates could have been originated from inhibition of StAR expression by collagen type IV.

Although testosterone production was increased in response to addition of hCG in ECM-exposed and unexposed cells and it was higher at all concentrations of hCG in the control than ECM group, it was shown that cells cultured on ECM-coated plates were more responsive to hCG compared to those cultured on uncoated plates. Gene expression analysis in the present study showed that ECM increased LH receptor expression, which probably contributed to greater responsiveness of Leydig cells cultured on ECM to hCG. Proteoglycans have been shown to modulate testosterone production by Leydig cells in rat.^{28,29} Factors neutralizing heparan sulfate proteoglycans have been observed to decrease testosterone production in LH-stimulated Leydig cells, suggesting that heparan

sulfate proteoglycans have positive impact on testosterone production through modulating LH receptor.²⁹ On the other hand, no significant effect of collagen type IV on the specific binding of hCG to LH receptor was observed in rat.³⁷ Accordingly, the greater responsiveness of Leydig cells to hCG in ECM than control group could have resulted from the modulating effect of heparan sulfate proteoglycan on LH receptor.

Indeed, different types of ECM are composed of a great variety of molecules with various functional and structural properties, which give each type of ECM specific capabilities for regulating the function and development of cells.¹⁸ In the present study, it was revealed that a combination of laminin, collagen type IV, heparan sulfate proteoglycan and entactin, which were derived from Engelbreth-Holm-Swarm mouse sarcoma, modulated factors contributing to testosterone synthesis and thus impacted testosterone production. Further studies are required to investigate the specific role of each ECM molecule as well as their interactive effect during the process of testosterone synthesis. Furthermore, the downstream mechanisms whereby ECM molecules influence testosterone production need to be elucidated.

In conclusion, the present study revealed that ECM decreased testosterone production by Leydig cells, through inhibiting StAR expression. Considering the permissive effect of ECM on SSCs differentiation, inhibition of testosterone production was probably one of the mechanisms whereby ECM facilitated SSCs differentiation. Additionally, it was shown that Leydig cells cultured on ECM were more responsive to hCG, which could have stemmed from the stimulatory effect of ECM molecules on LH receptor expression.

Acknowledgments

The present study was supported by Center of Excellence in Application of Stem cells in Cell Therapy and Tissue Engineering, Faculty of Veterinary Medicine, University of Tehran and Theriogenology Association, Faculty of Veterinary Medicine, University of Tehran. The authors would like to thank Dr. Shiva Shafiei, Dr. Zohreh Mazaheri and Mr. Shahram Pour Beiranvand for their kind assistance in the present study.

References

1. Oatley JM, Brinster RL. Regulation of spermatogonial stem cell self-renewal in mammals. *Annu Rev Cell Dev Biol* 2008; 24: 263-286.
2. Tseng YT, Liao HF, Yu CY, et al. Epigenetic factors in the regulation of prospermatogonia and spermatogonial stem cells. *Reproduction* 2015; 150: 77-91.
3. McLachlan RI, O'Donnell L, Meachem SJ, et al. Identification of specific sites of hormonal regulation

- in spermatogenesis in rats, monkeys and man. *Recent Prog Horm Res* 2002; 57: 149-179.
4. Walker WH. Testosterone signaling and the regulation of spermatogenesis. *Spermatogenesis* 2011; 1: 116-120.
 5. Andersson AM, Jørgensen N, Frydelund-Larsen L, et al. Impaired Leydig cell function in infertile men: A study of 357 idiopathic infertile men and 318 proven fertile controls. *J Clin Endocrinol Metab* 2004; 89: 3161-3167.
 6. Rich KA, De Kretser DM. Effect of differing degrees of destruction of the rat seminiferous epithelium on levels of serum follicle stimulating hormone and androgen binding protein. *Endocrinology* 1977; 101: 959-968.
 7. Rich KA, Kerr JB, de Kretser DM. Evidence for Leydig cell dysfunction in rats with seminiferous tubule damage. *Mol Cell Endocrinol* 1979; 13: 123-135.
 8. Singh J, O'Neill C, Handelsman DJ. Induction of spermatogenesis by androgens in gonadotropin-deficient (hpg) mice. *Endocrinology* 1995; 136: 5311-5321.
 9. Meng J, Holdcraft RW, Shima JE, et al. Androgens regulate the permeability of the blood-testis barrier. *Proc Natl Acad Sci USA* 2005; 102: 16696-16700.
 10. Meng J, Greenlee AR, Taub CJ, et al. Sertoli cell-specific deletion of the androgen receptor compromises testicular immune privilege in mice. *Biol Reprod* 2011; 85: 254-260.
 11. Wang RS, Yeh S, Chen LM, et al. Androgen receptor in Sertoli cell is essential for germ cell nursery and junctional complex formation in mouse testes. *Endocrinology* 2006; 147: 5624-5633.
 12. Willems A, Batlouni SR, Esnal A, et al. Selective ablation of the androgen receptor in mouse Sertoli cells affects Sertoli cell maturation, barrier formation and cytoskeletal development. *PLoS One* 2010; 5: e14168.
 13. Holdcraft RW, Braun RE. Androgen receptor function is required in Sertoli cells for the terminal differentiation of haploid spermatids. *Development* 2004; 131: 459-467.
 14. O'Donnell L, McLachlan RI, Wreford NG, et al. Testosterone promotes the conversion of round spermatids between stages VII and VIII of the rat spermatogenic cycle. *Endocrinology* 1994; 135: 2608-2614.
 15. O'Donnell L, McLachlan RI, Wreford NG, et al. Testosterone withdrawal promotes stage-specific detachment of round spermatids from the rat seminiferous epithelium. *Biol Reprod* 1996; 55: 895-901.
 16. Payne AH. Hormonal regulation of cytochrome P450 enzymes, cholesterol side-chain cleavage and 17 alpha-hydroxylase/C17-20 lyase in Leydig cells. *Biol Reprod* 1990; 42: 399-404.
 17. Stocco DM, Clark BJ. Regulation of the acute production of steroids in steroidogenic cells. *Endocr Rev* 1996; 17: 221-244.
 18. Rosso F, Giordano A, Barbarisi M, et al. From cell-ECM interactions to tissue engineering. *J Cell Physiol* 2004; 199: 174-180.
 19. Schwartz MA, Baron V. Interaction between mitogenic stimuli, or, a thousand and one connections. *Curr Opin Biol* 1999; 11: 197-202.
 20. Taipale J, Keski-Oja J. Growth factors in the extracellular matrix. *FASEB J* 1997; 11: 51-59.
 21. Abbott A. Cell culture: Biology's new dimension. *Nature* 2003; 424: 870-872.
 22. Zhang S. Beyond the petri dish. *Nat Biotechnol* 2004; 22: 151-152.
 23. Akbarinejad V, Tajik P, Movahedin M, et al. Effect of extracellular matrix on bovine spermatogonial stem cells and gene expression of niche factors regulating their development *in vitro*. *Anim Reprod Sci* 2015; 157: 95-102.
 24. Lee JH, Kim HJ, Kim H, et al. *In vitro* spermatogenesis by three-dimensional culture of rat testicular cells in collagen gel matrix. *Biomaterials* 2006; 27: 2845-2853.
 25. Lee JH, Gye MC, Choi KW, et al. *In vitro* differentiation of germ cells from nonobstructive azoospermic patients using three-dimensional culture in a collagen gel matrix. *Fertil Steril* 2007; 87: 824-833.
 26. Vernon RB, Lane TF, Angello JC, et al. Adhesion, shape, proliferation, and gene expression of mouse Leydig cells are influenced by extracellular matrix *in vitro*. *Biol Reprod* 1991; 44: 157-170.
 27. Diaz ES, Pellizzari E, Meroni S, et al. Effect of extracellular matrix proteins on *in vitro* testosterone production by rat Leydig cells. *Mol Reprod Dev* 2002; 61: 493-503.
 28. Grudet N, Bonnamy PJ, Le Goff D, et al. Role of proteoglycans on testosterone synthesis by purified Leydig cells from immature and mature rats. *J Steroid Biochem Mol Biol* 1999; 68: 153-162.
 29. McFarlane JR, Laslett A, de Kretser DM, et al. Evidence that heparin binding autocrine factors modulate testosterone production by the adult rat Leydig cell. *Mol Cell Endocrinol* 1996; 118: 57-63.
 30. Mossa F, Jimenez-Krassel F, Folger JK, et al. Evidence that high variation in antral follicle count during follicular waves is linked to alterations in ovarian androgen production in cattle. *Reproduction* 2010; 140: 713-720.
 31. Miller WL. Steroidogenic acute regulatory protein (StAR), a novel mitochondrial cholesterol transporter. *Biochim Biophys Acta* 2007; 1771: 663-676.
 32. Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2^{-ΔΔC_T} method. *Methods* 2001; 25: 402-408.
 33. Littell RC, Henry PR, Ammerman CB. Statistical analysis of repeated measures data using SAS procedures. *J Anim Sci* 1998; 76: 1216-1231.
 34. Shetty G, Wilson G, Huhtaniemi I, et al. Testosterone inhibits spermatogonial differentiation in juvenile spermatogonial depletion mice. *Endocrinology* 2001; 142: 2789-2795.

35. Shetty G, Weng CC, Porter KL, et al. Spermatogonial differentiation in juvenile spermatogonial depletion (jsd) mice with androgen receptor or follicle-stimulating hormone mutations. *Endocrinology* 2006; 147: 3563-3570.
36. Zhou W, Wang G, Small CL, et al. Gene expression alterations by conditional knockout of androgen receptor in adult Sertoli cells of Utp14b jsd/jsd (jsd) mice. *Biol Reprod* 2011; 84: 400-408.
37. Diaz ES, Pellizzari E, Casanova M, et al. Type IV collagen induces down-regulation of steroidogenic response to gonadotropins in adult rat Leydig cells involving mitogen-activated protein kinase. *Mol Reprod Dev* 2005; 72: 208-215.