ORIGINAL ARTICLE

Veterinary Research Forum. 2020; 11 (3) 243 - 248 doi: 10.30466/vrf.2018.90574.2192

Veterinary Research Forum

Journal Homepage: vrf.iranjournals.ir

Molecular analysis of pyrethroid resistance in *Cimex hemipterus* (Hemiptera: Cimicidae) collected from different parts of Iran

Awat Samiei¹, Mousa Tavassoli^{1*}, Karim Mardani²

¹ Department of Pathobiology, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran; ² Department of Food Hygiene and Quality Control, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran.

Article Info	Abstract
Article history:	The present study was aimed to assess the bedbugs susceptibility to pyrethroid
	insecticides using molecular analysis. With the aid of pest control companies, adult bedbugs
Received: 23 July 2018	were collected from various places such as hotels, residential houses, and industrial buildings
Accepted: 18 September 2018	in seven cities highly crowded with domestic and foreign tourists in Iran from May 2016 to
Available online: 15 September 2020	August 2017. Bedbugs were colonized in the laboratory to evaluate their resistance to
	pyrethroid using insecticide resistance bioassay. Genomic DNA was extracted from
Keywords:	susceptible and resistant bedbugs. At first, specie specific primers targeting cytochrome
-	oxidase subunit I (COI) gene was performed to confirm Cimex hemipterus species. Then, kdr-
Bedbugs	like gene was examined for point mutation using PCR and nucleotide sequencing. Bioassay
Cimex hemipterus	showed that 11 out of 35 examined bedbugs were resistant to pyrethroids (31.43%; 95.00%
Insecticide	confidence interval: 29.48-33.08%). The DNA sequencing showed that all examined bedbugs
Pyrethroids	collected from Tehran province had homozygous V419L kdr-like gene mutations. The level of
Resistance	pyrethroid resistance found in the collected bugs from Tehran province indicated that this
	phenomenon has already been prevailed in the site and prompts the need to reevaluate the
	large use of pyrethroids to control the bedbugs.

© 2020 Urmia University. All rights reserved.

Introduction

Two common and the most important species of Cimicidae family, members of cimicids are Cimex lectularius and *Cimex hemipterus*, known as bedbugs. The usual hosts for these blood-feeding ectoparasites are bats, birds, and particularly humans.1 It has been well documented that bedbugs harbor at least 40 human pathogens, however, there is no tangible evidence regarding their ability to transmit these pathogens to human.¹⁻⁵ There are physical trauma, wheals (typically 5.00-20.00 mm), and itch from the bug's bites,⁵ and the severity of the reaction depends on the individual's sensitivity.6 Skin lesions in the bug infestations may include dermatitis, ecthyma, folliculitis, impetigo, and lymphangitis.⁶ Some people suffer from allergic reactions due to the bedbug saliva, causing red inflammation and triggering anaphylaxis in certain cases.⁶ There are also Reduviid bugs (Hemiptera: Reduviidae), including the bedbug distant cousin Triatoma infestans, also called the kissing bug, one of the species which

spreads Chagas disease from the southern United States down through South America by passing the tropical parasite Trypanosoma cruzi to hosts as they sleep,² emphasizing that this family does not exist in Iran.

Cimex hemipterus, is a wingless, red-brown, bloodsucking insect with a maximum length of eight mm and a lifespan ranging from four months up to one year. All the daytime, bedbugs disappear and hide in inaccessible places such as cracks and crevices in beds, wooden furniture, floors, and walls and re-appear at night to feed on their preferred host, humans.7

A resurgence of bedbugs has been reported from the United States, Canada, Australia, Europe, and some Asian countries during the past 15 years.⁸⁻¹⁴ The reasons for the explosion have not yet been clarified, however, several factors such as increased rate of international travel, reduced use of residual insecticides indoors, and insecticide resistance may play a role.15

Recent reports are claiming that several species of arthropod pests have developed considerable resistance to

*Correspondence:

Mousa Tavassoli. DVM, PhD

Department of Pathobiology, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran E-mail: m.tavassoli@urmia.ac.ir



This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License which allows users to read, copy, distribute and make derivative works for non-commercial purposes from the material, as long as the author of the original work is cited properly.

available insecticides, posing great danger to human life and needing immediate attention.¹⁶⁻²⁰ Mechanisms beyond the developed resistance in bedbugs against the pyrethroids, the most commonly used pesticide, including behavioral resistance,²¹ cuticular resistance,²² metabolic resistance,23 and target-site insensitivity resistance are because of the substitute mutations at the DNA level.²⁴ The most prominent mechanism by which several species of insects develop resistance to insecticides is mutations occurred in the voltage-gated sodium channel (VGSC) reducing the affinity of the receptor.^{20,25} Target site insensitivity of this type is known as 'knockdown' (or kdrtype) resistance, named firstly after a phenomenon observed in houseflies survived following being knocked down by DDT and subsequently demonstrating crossresistance to pyrethrins.^{26,27} The *kdr*-type target-site insensitivity resistance has been extensively studied in both C. lectularius and C. hemipterus, it is present everywhere within pyrethroid-resistant field-strains and plays a significant role in the expression of pyrethroid resistance. V419L and L925I mutations in a VGSC asubunit gene of C. lectularius, have also been used as a molecular marker of resistance to pyrethroids in a few studies.^{24,28,29} To date there is no study underlying the resistant mechanisms of bedbugs against pyrethroids in Iran. Therefore, the current study was aimed to assess the susceptibility of bedbugs collected from different parts of Iran to pyrethroid insecticides at the molecular level.

Materials and Methods

Regions and sample (bedbug) collection. There are four different geographical zones in Iran, known as region 1: Caspian Sea (temperature: 8.00 - 26.00 °C, annual rainfall: 400 - 1500 mm), region 2: Mountainous area (temperature: -5.00 - 29.00 °C, annual rainfall: 200-500 mm), region 3: Persian Gulf (temperature: 12.60 - 35.00 °C, annual rainfall: 200-300 mm), and region 4: The Central Desert (temperature: -4.00 - 44.00 °C, rainfall: less than 100 mm). The pattern of bedbug species distribution for the four different areas was determined according to the method of Skerman and Hillard.³⁰

With the aid of pest control companies, adult bedbugs were collected from various places in each region such as hotels, residential houses, and industrial buildings before treating the infested areas with insecticide from May 2016 to August 2017. From each of the regions one or two cities were selected: Region 1: Sari (36.5659° N, 53.0586° E) and Rasht (37.2682° N, 49.5891° E), region 2: Tehran (35.6892° N, 51.3890° E) and Hamadan (34.7989° N, 48.5150° E), region 3: Ahvaz (31.3183° N, 48.6706° E) and Bandar Abbas (27.1832° N, 56.2666° E), and region 4: Semnan (35.2256° N, 54.4342° E).

Bedbug samples were transferred to the parasitology laboratory, Faculty of Veterinary Medicine, Urmia University for identification under a stereomicroscope (Olympus SZ61; Olympus Corporation, Tokyo, Japan), using morphological features and the taxonomic key of Usinger. Useful diagnostic features included rostrum length, mesothoracic coxae separation, antennal segment length, as well as pronotal anterior margin and hairs.¹

Insecticide resistance bioassay. The resistance of the collected bedbugs from four regions were examined using the World Health Organization (WHO) insecticideimpregnated papers. Each paper (8.50 cm diameter) was impregnated with 500 µL of 10.00% cypermethrin (100 ppm; National Agrochemical Co., Tehran, Iran) according to the recommended dosage of the manufacturer. Each insecticide-impregnated paper was cut into 3.00 × 5.00 cm pieces, rolled, and placed inside a glass test tube (length × internal diameter: 12.50 × 1.50 cm). A piece of clean paper $(3.00 \times 5.00 \text{ cm})$ impregnated with isotonic sodium chloride solution was used as a control group similar to the impregnated paper was rolled and placed inside another glass test tube. Alive and motile adult bedbugs of either sex from each region (five samples per city) were carefully transferred into each tube and closed with finemesh gauze. The tubes were set upright in the holding rack and placed in a plastic box $(20.00 \times 20.00 \times 15.00 \text{ cm})$ lined inside with damp cloths to obviate very low humidity. The box was closed for the appropriate exposure period. Bedbugs were maintained in an incubator at 24.00 ± 2.00 °C and 55.00 - 65.00% relative humidity. The concentrations of insecticides and exposure times used were recommended by WHO.31 Mortality rates of the bedbugs exposed to cypermethrin were recorded after 24 hr of exposure to insecticide or isotonic chloride solution. Bedbugs were considered as dead if there were no vital signs or minor vital signs present, merely internal gut movements, movements of antennae, minimal leg movements with or without stimulation by a forceps. Each bioassay was replicated three times against each colony of bedbugs.

DNA extraction. Genomic DNA was extracted from individual tropical bedbug of each collection site using a DNA extraction kit (MBST, Tehran, Iran) following the manufacturer's instructions and as per the method described by Shayan and Rahbari.³² Briefly, bug samples were ground by pestle and placed into 1.50 mL microcentrifuge tube and DNA extraction was performed according to the kit's protocol. Total DNA was eluted in 100 μ L of elution buffer and its quality and concentration was determined using the NanoDrop (2000c; Thermo Fisher Scientific, Waltham, USA) and stored at – 20.00 °C for further procedures.

Amplification of cytochrome oxidase subunit I (COI) gene. Amplification of a 685 bp fragment of the COI gene using species-specific PCR primers LEP-F (5'-ATTCA ACCAATCATAAAGATATNG G- 3') and LEP-R (5'-TAWACT TCWGGRTGTCCRAARAAT CA-3') described by Balvin *et al* was performed to confirm *C. hemipterus* species.³³

Genotyping of kdr-like gene. The PCR amplification of two 474-bp and 744-bp amplicons of the VGSC subunit gene (kdr-like gene) was performed using primers (forward KDR1: 5'AACCTGGATATACATGCCTTCAAGG3' and reverse KDR1: 5_TGATGGAGATTTTGCCACTGATG3 for fragment 1; forward KDR2: 5_GGAATTGAAGCTGCCAT GAAGTTG3 and reverse KDR2: 5_TGCCTATTCTGTCGAAA GCCTCAG3 for fragment 2) described previously.³⁴ PCR was carried out in 25.00 µL reaction volume consisting of 2.50 µL 10X PCR buffer, 2.00 µL MgCl₂ (50.00 mM), 0.50 µL dNTPs (10.00 mM), 3.00 µL DNA template, 10.00 pmol of each forward and reverse primer (0.50 µL for each) for either V419L and L925I, 0.50 µL Taq DNA polymerase (SinaClon, Tehran, Iran) and 15.50 µL ddH₂O. PCR cycling conditions were as follows: Initial denaturation at 94.00 °C for 10 min followed by 40 cycles of 94.00 °C for 40 sec, 52.00 °C (V419L) and 55.00 °C (L925I) for 40 sec, 72.00 °C for 40 sec and a final extension step of 72.00 °C for 10 min. PCR products were visualized on 1.50% agarose gel stained with safe DNA dye under UV-Transilluminator (20M; BTS, Tokyo, Japan).

The PCR products were sent to SinaClon Company (Tehran, Iran) for purification and nucleotide sequencing. The obtained sequences were compared to the wild-type sequences retrieved from GenBank (GU123927 and GU123928). The presence of single nucleotide polymorphisms (V419L and L9251) was confirmed by reading both forward and reverse strands.

Statistical analysis. The obtained data used for mortality bioassay was analyzed using the SPSS program (version 22.0; IBM Corp., Armonk, USA).

Results

Mortality bioassay. Only alive and motile adult bedbugs were selected for the test. In total, 40 bedbugs were subjected individually to the test. Five bedbugs were used as controls, of which none was dead at the end of the test. Among the 35 bedbugs exposed to insecticide, 11 (31.43%) were reported as resistant (95.00% confidence interval: 29.48-33.08%) and 24 (68.57%) were susceptible (95.00% confidence interval: 67.82-71.62%). The Most resistant bedbugs (5 out 11) were collected in two highly infested places located in Tehran province.

Genotyping of *kdr***-like gene.** At first, amplification of the COI gene for bedbugs showed a consistent amplification with strong bands that specify all samples belong to *C. hemipterus*. Subsequently, the DNA of all 35 bedbugs exposed to insecticide during the mortality bioassay was extracted. Both parts of the *kdr*-like gene containing codons 419 and 925 were successfully amplified and sequenced for 20 bedbugs (Fig. 1), including nine susceptible and 11 resistant bedbugs according to the bioassays. In *C. hemipterus* field populations (Table 1), all sequenced bedbugs had homozygous wild-type L925I

codon and homozygous mutated V419L codon related to Tehran province of *kdr*-like gene (Fig. 2). In other words, the existence of resistance alleles at the V419L mutation for all populations was 0.0 except for Tehran.

GU123927	GTCCGTGGCACAT GTTGTTCTTCATAGTAATTATATTTCCTGGGATCATTCTACCTCGTCA	60
THE-70	GTCCGTGGCACATGTTGTTCTTCATAGTAATTATATTCCTGGGATCATTCTACCTCCTCA	60
GU123927	ACTTGATTTTAGCTATTGTCGCTATGTCATATGATGAACTGCAAAAGAAGGCGGAAGAAG	120
THE-70	ÅĊTTGAŤŤŤŤĂGCTATTGTCGCTATGTCATATGATGAACTGCAAAAGAAGGCGGAAGAAG	120
GU123927	AAGAAGCTGCTGAAGAGGAAGCCCTTAGGGTATGCTTTAGCAAGGAAGCAATTGAAAATA	180
THE-70	AÁĠAAGCTGCTGÁAGAGGAAGCCCTTÁGGGTATGCTTTAGCAAGGAAGCAATTGAAAATA	180
GU123927	TCTTGAATCACATGGGAATTCAAAATATTTAGGAGGCAGAGGAAGCAGCTCAAGCTAAGG	240
THE-70	TCTTGAATCACATGGGAATTCAAAATATTTAGGAGGCAGAGGAAGCAGCTCAAGCTAAGG	240

Fig. 1. Alignment of a part of the sequence of the *kdr*-like gene, encompassing codon 419 (in red), from a pyrethroid-susceptible bedbug [Genbank access number: GU123927] and adult bedbugs collected in the study.

Table 1. The *kdr*-type gene mutations found in *C. hemipterus* examined in the present study from May 2016 to August 2017.

Cities	V419L	L925I
Sari	-	-
Rasht	-	-
Hamadan	-	-
Ahvaz	-	-
Bandar Abbas	-	-
Semnan	-	-
Tehran	*	-

* Homozygous mutated V419L codon related to Tehran city of the *kdr*-like gene.



Fig. 2. Chromatograms showing differences between *kdr*-type gene sequences of pyrethroid susceptible and resistant bedbugs. On the left-hand side, the VGSC amino acid substitution that results in a kdr mutation is indicated, and the corresponding base pair change is bolded and underlined. The same base pairs are indicated by an arrow in the chromatogram sequence, and its identity is boxed on the chromatogram base-pair readout.

Discussion

Bedbugs were a common pest up to the mid-20th century, however, with the advent of synthetic insecticides, their population considerably was declined particularly in the developed countries.³⁵ On the other hand, during the past two decades a resurgence of bedbugs infestations have been reported simultaneously

all around the world,⁵ including Europe,^{36,37} Africa,³⁸ South America,³⁹ Asia,^{40,41} Australia,⁴² and North America.⁴³ It is estimated that the resurgence of bedbugs has incurred severe economic losses which cost billions of dollars,¹¹ including 200 million AUD in Australia since the resurgence started in 2011.⁴⁴ The incurred losses were due to eradication charges, lost revenue, and discarded infested items.⁴⁵ It has been claimed that a blend of various factors such as insecticide resistance, increased international travel, and poor control practices probably have led to the resurgence of bedbugs.²⁵

Bioassays determined that pyrethroid-resistant bedbugs were present in several cities in Iran. PCO (pest control operators) usually uses pyrethroid insecticides in different regions of Iran. They use mostly 10.00% cypermethrin). People frequently use these products, which are easily available in local stores. As a result, pyrethroid insecticides are often improperly used in terms of concentrations and the number of applications, which may induce unnecessary insecticide pressure on bedbugs and the other pests.

Some bedbugs examined in the present study showed a point mutation in V419L location leading to the resistance to pyrethroids. Resistance to pyrethroids was not always associated with the V419L mutation in bedbugs in the USA and this mutation was not always associated with the L925I mutation to form a resistant haplotype.^{24,28} The present data is the only results of genotyping of C. hemipterus based on a kdr-like gene in Iran. The presence of *kdr*-type mutations was detected for the first time in *C*. lectularius collected in New York, USA, using two target sites in the kdr-like gene for point mutations (V419L and L925I).²⁴ Then a high rate of the mutation. almost 88.00% was detected in 110 field-strains of C. lectularius collected across the USA.23 Furthermore, researches have subsequently disclosed the presence of mutations in *kdr*-type genes in parathyroid-resistant bedbugs around the world. For example, in France, all specimens of C. lectularius collected from a single multi-occupancy high-rise apartment complex were reported to homogeneously have only haplotype B (L925I) kdr-type mutations.³³ Besides, time changing mutations of both V419L and L925I were revealed in *C. lectularius* in Korea²⁸. The *kdr*-type gene analysis of C. hemipterus strains determined that the V419L, L925I, and I936F mutations were absent from the specimens investigated, although two other mutations, M918I and L1014F, were putative kdr-type mutations as they appear in other pyrethroid-resistant insects.⁴⁶

Our results showed that pyrethroids resistant bedbugs were common in many places of Tehran, since PCOs mostly use pyrethroid insecticides in this city and its suburbs. Also, due to heavily populated and passenger traffic, people often use pyrethroid insecticides which are easily available in the local stores. Although studies are limited, to date all investigations examining *kdr*-type genes have revealed a high proportion of the mutations in this gene suggesting that these mutations are probably widespread in bedbugs across the world. The continuous use of pyrethroid insecticides, despite not being very efficacious, might be contributed to the maintenance of high frequency of *kdr*-type gene mutations.

The evolutionary mechanisms beyond the respective *kdr*-type gene mutations among bedbug species have not been completely understood yet and this must be clarified in future studies. Furthermore, it is important that pest control managers tightly adhere to "best practice" as defined in the various bedbug management industry standards to reduce the risk of treatment failure. In conclusion, the current study was the first effort to investigate the distribution and extent of *kdr*-type gene mutations in bedbug populations across Iran. Also, the results obtained from this study suggested that the bedbugs in Tehran province have developed resistance to various groups of pyrethroid insecticides which are the most common insecticides used for pest control.

Acknowledgments

The authors would like to thank Iranian Pest Control Companies for their cooperation in collecting bedbugs. We are also grateful to Dr. Ali Nazarizadeh and Dr. Behnaz Ghorbanzadeh for their assistance in manuscript preparation and helpful advice.

Conflict of interest

The authors declare no potential conflict of interest.

References

- 1. Usinger RL. Monograph of cimicidae (Hemiptera-Heteroptera). Entomological Society of America. College Park, USA: Thomas Say Foundation 1966; 1.
- Cooper RA, Harlan H. Ectoparasites: Bed bugs and kissing bugs. In: Hedges S (Ed). Mallis handbook of pest control. 9th ed. Cleveland, USA: GIE Publishing 2004; 494-529.
- Goddard J, de Shazo R. Bed bugs (*Cimex lectularius*) and clinical consequences of their bites. J Am Med Assoc 2009; 301(13): 1358-1366.
- 4. Wang C, Tsai WT, Cooper R, et al. Effectiveness of bed bug monitors for detecting and trapping bed bugs in apartments. J Econ Entomol 2011; 104(1): 274-278.
- 5. Doggett SL, Dwyer DE, Peñas PF, et al. Bed bugs: Clinical relevance and control options. Clin Microbiol Rev 2012; 25(1): 164-192.
- 6. Zorrilla-Vaca A, Silva-Medina, MM, Escandón-Vargas, K. Bed bugs, *Cimex* spp.: Their current world resurgence and healthcare impact. Asian Pac J Trop Dis 2015; 5(5): 342-352.

- Elston DM, Stockwell S. What's eating you? Bedbugs. Cutis 2000; 65(5): 262-264.
- 8. Hwang SW, Svoboda TJ, De Jong IJ, et al. Bed bug infestations in an urban environment. Emerg Infect Dis 2005; 11(4): 533-538.
- 9. Cooper R. Bed bugs-Still more questions than answers: A need for research and public awareness. Am Entomol 2006; 52(2): 111-112.
- 10. Romero A, Potter MF, Potter DA et al. Insecticide resistance in the bed bug: A factor in the pest's sudden resurgence? J Med Entomol2007; 44(2): 175-178.
- 11. Doggett SL, Russell RC. The resurgence of bed bugs, Cimex spp. (Hemiptera: Cimicidae) in Australia. In proceedings: 6th International conference on urban pests. Budapest, Hungary. 2008; 6: 407-425.
- 12. Kilpinen O, Jensen KMV, Kristensen M. Bed bug problems in Denmark, with a European perspective. In proceedings: 6th International conference on urban pests. Budapest, Hungary 2008; 13-16
- 13. Wang L, Xu Y, Zeng L. Resurgence of bed bugs (Hemiptera: Cimicidae) in mainland China. Fla Entomol 2013; 96(1): 131-136.
- 14. Zulaikha Z, Hafiz ABA, Hafis ARA, et al. A survey on the infestation levels of tropical bed bugs in Peninsular Malaysia: Current updates and status on resurgence of Cimex hemipterus (Hemiptera: Cimicidae). Asian Pac J Trop Dis 2016; 6(1): 40-45.
- 15. Motevalli Haghi SF, Behbodi M, Hajati H, et al. Prevalence of bed bug (*Cimex lectularius*) in human settlement area of Bahnamir, Iran. Asian Pac J Trop Dis 2014; 4 (Suppl. 2): S786-S789.
- 16. Hemingway J, Field L, Vontas J. An overview of insecticide resistance. Science 2002; 298(5591):96-97.
- 17. Hemingway J, Ranson H. Insecticide resistance in insect vectors of human disease. Annu Rev Entomol 2000; 45(1):371-391.
- Horowitz A, Denholm I. Impact of insecticide resistance mechanisms on management strategies. In: Ishaaya I. (Ed). Biochemical sites of insecticide action and resistance. Berlin, Germany: Springer-Verlag 2001; 323-338.
- 19. Roush R, Tabashnik BE. Pesticide resistance in arthropods: Berlin, Germany: Springer Science & Business Media 2012; 1-4.
- 20. Yu SJ. Insecticide resistance. The Toxicology and Biochemistry of Insecticides. 2nd ed. Boca Raton, USA: CRC Press 2015; 257-321.
- 21. Romero A, Potter MF, Haynes KF. Behavioral responses of the bed bug to insecticide residues. J Med Entomol 2009; 46(1): 51-57.
- 22. Koganemaru R, Miller DM, Adelman ZN. Robust cuticular penetration resistance in the common bed bug (*Cimex lectularius* L.) correlates with increased steady-state transcript levels of CPR-type cuticle protein genes. Pest Biochem Physiol 2013; 106(3): 190-197.

- 23. Zhu F, Gujar H, Gordon JFR et al. Bed bugs evolved unique adaptive strategy to resist pyrethroid insecticides. Sci Rep 2013; 3:1456. doi:10.1038/ srep01456.
- 24. Yoon KS, Kwon DH, Strycharz JP, et al. Biochemical and molecular analysis of deltamethrin resistance in the common bed bug (Hemiptera: Cimicidae). J Med Entomol 2008; 45(6): 1092-1101.
- 25. Davies TG, Field LM, Williamson MS. The re-emergence of the bed bug as a nuisance pest: implications of resistance to the pyrethroid insecticides. Med Vet Entomol 2012; 26(3): 241-254.
- 26. Busvine JR. Mechanism of resistance to insecticide in houseflies. Nature 1951; 168(4266):193-195.
- 27. Soderlund DM. Pyrethroids, knockdown resistance and sodium channels. Pest Manag Sci 2008; 64(6): 610-616.
- 28. Seong KM, Lee DY, Yoon KS, et al. Establishment of quantitative sequencing and filter contact vial bioassay for monitoring pyrethroid resistance in the common bed bug, *Cimex lectularius*. J Med Entomol 2010; 47(4): 592-599.
- 29. Zhu F, Wigginton J, Romero A, et al. Widespread distribution of knockdown resistance mutations in the bed bug, *Cimex lectularius* (Hemiptera: Cimicidae), populations in the United States. Arch Insect Biochem Physiol 2010; 73(4): 245-257.
- 30. Skerman K, Hillard J. A handbook for studies of helminth parasites of ruminants. Near East Animal Health Institutes, Iran Unit, United Nations Development Programme/Special fund 1967; 1-3.
- 31. WHO. World Health Organization. Instructions for determining the susceptibility or resistance of adult bedbugs to insecticides 1981; WHO/VBC/81.809.
- 32. Shayan P, Rahbari S. Simultaneous differentiation between Theileria spp. and Babesia spp. on stained blood smear using PCR. Parasitol Res 2005; 97(4): 281-286.
- 33. Balvín O, Munclinger P, Kratochvíl L, et al. Mitochondrial DNA and morphology show independent evolutionary histories of bedbug *Cimex lectularius* (Heteroptera: Cimicidae) on bats and humans. Parasitol Res 2012; 111(1): 457-469.
- 34. Durand R, Cannet A, Berdjane Z, et al. Infestation by pyrethroid resistance bed bugs in the suburb of Paris, France. Parasite 2012; 19(4): 381-387.
- 35. Potter MF. The history of bed bug management-with lessons from the past. Am Entomol 2011; 57(1): 14-25.
- 36. Sadílek Da, Šťáhlavský F, Vilímová J, et al. Extensive fragmentation of the X chromosome in the bed bug *Cimex lectularius* Linnaeus, 1758 (Heteroptera, Cimicidae): A survey across Europe. Comp Cytogenet 2013, 7(4): 253-269.
- 37. Vander Pan A, Kuhn C, Schmolz E, et al. Studies on pyrethroid resistance in *Cimex lectularius* (Hemiptera:

Cimicidae), in Berlin, Germany. In proceedings: 8th International conference on urban pests. Zurich, Switzerland. 2014; 89-95.

- 38. Omudu EA, Kuse CN. Bedbug infestation and its control practices in Gbajimba: A rural settlement in Benue state, Nigeria. J Vector Borne Dis 2010; 47(4): 222-227.
- Posso CE, Wolff MI, de Ulloa PC. The bed bugs *Cimex lectularius*: Urban problem in the Andean region of Colombia. In proceedings: 7th International conference on urban pests. São Paulo, Brazil 2011; 371.
- 40. Lee IY, Ree HI, An SJ, et al. Reemergence of the bedbug *Cimex lectularius* in Seoul, Korea. Korean J Parasitol 2008; 46(4): 269-271.
- 41. Tawatsin A, Thavara U, Chompoosri J, et al. Insecticide resistance in bedbugs in Thailand and laboratory evaluation of insecticides for the control of *Cimex hemipterus* and *Cimex lectularius* (Hemiptera: Cimicidae).

J Med Entomol 2011; 48(5): 1023-1030.

- 42. Doggett SL, Geary MJ, Crowe WJ, et al. Has the tropical bed bug, *Cimex hemipterus* (Hemiptera: Cimicidae), invaded Australia? Environ Health 2003; 3(4): 80-82.
- 43. Krueger L. Don't get bitten by the resurgence of bed bugs- properly identifying a bed bug infestation is the key to quick control. Pest Control 2000; 28(3): 58-64.
- 44. Doggett SL, Orton CJ, Lilly DG, et al. Bed bugs: The Australian response. Insects 2011; 2(2): 96-111.
- 45. Schaefer CW. Bed bugs (Cimicidae). In: Schaefer CW, Panizzi AR (Eds). Heteroptera of economic importance. Boca Raton, USA: CRC Press 2000; 519-538.
- 46. Dang K, Toi CS, Lilly DG, et al. Identification of putative *kdr* mutations in the tropical bed bug, *Cimex hemipterus* (Hemiptera: Cimicidae). Pest Manag Sci 2015; 71(7):1015-1020.