



1st International Congress of Veterinary Medicinal Plants and Traditional Medicine



Plant Products and Pharmacology
Effects of Plant Products on Animal and Zoonotic Pathogens
Plant Products and Food Science
Traditional Veterinary Medicine
Phytochemistry and Biotechnology in Medicinal Plants

Topics:

October
21 & 22
2024
Urmia - Iran



www.licvmp.urmia.ac.ir
www.licvmp.com

Probiotic potential of autochthonous *Lactobacillus* species from buffalo calves in controlling multidrug resistant *Escherichia coli*

Srivani Moturi^{1*}, Lakshmi Kavitha Kommalapati², Muralidhar Metta³, Venkata Seshiah Chappidi⁴, Srilakshmi Jatavathu⁵

¹ Department of Veterinary Microbiology, N.T.R College of Veterinary Science, Sri Venkateswara Veterinary University, Vijayawada, India; ² Department of Veterinary Microbiology, College of Veterinary Science, Sri Venkateswara Veterinary University, Tirupati, India; ³ Department of Animal Genetics and Breeding, College of Veterinary Science, Sri Venkateswara Veterinary University, Vizianagaram, India; ⁴ Livestock Farm Complex, N.T.R College of Veterinary Science, Sri Venkateswara Veterinary University, Vijayawada, India; ⁵ Junior Research Fellow, Department of Veterinary Microbiology, N.T.R College of Veterinary Science, Sri Venkateswara Veterinary University, Vijayawada, India.

Article Info	Abstract
Article history: Received: 29 September 2023 Accepted: 08 January 2024 Available online: 15 June 2024	The aim of this study was to investigate the probiotic potential of autochthonous <i>Lactobacillus</i> species isolated from buffalo calves against multidrug-resistant <i>Escherichia coli</i> . A total of 252 rectal swabs were collected from healthy neonatal buffalo calves under 30 days old from six districts of Andhra Pradesh, India in a completely randomized design from August 2019 to August 2021, of which 190 <i>Lactobacillus</i> strains were isolated based on cultural, morphological, biochemical and molecular tests. Out of 190 isolates, 57 showed high levels of auto-aggregation (> 80.00%) and hydrophobicity (> 60.00%) and 51 of the 57 isolates had a zone of inhibition greater than 15.00 mm in diameter against multidrug-resistant <i>E. coli</i> in an Agar well diffusion assay. Among the 51 isolates, 36 were found to be acid and bile tolerant and showed varying levels of sensitivity to antibiotics such as erythromycin, clindamycin, tetracycline, chloramphenicol, and ampicillin. Among the 36 isolates, <i>Limosilactobacillus reuteri</i> 178, <i>L. reuteri</i> 209, <i>L. fermentum</i> 182, <i>L. fermentum</i> 211, and <i>Lactiplantibacillus plantarum</i> 34 were non-hemolytic, and none of the isolates were able to hydrolyse gelatine. Therefore, these five autochthonous <i>Lactobacillus</i> species may be used in probiotic or synbiotic formulations against multidrug resistant <i>E. coli</i> in buffalo calves.
Keywords: Autochthonous probiotic Buffalo calves Multidrug resistant <i>E. coli</i>	

© 2024 Urmia University. All rights reserved.

Introduction

The natural microbiota of the digestive tract impacts on the biochemistry, immunology, physiology, and non-specific host resistance to infectious diseases.¹ In neonatal calves, stress, dietary changes, and environmental factors contribute to the colonization of opportunistic pathogens in the gut, leading to digestive problems and diarrhea. Antibiotics are frequently used to treat calf's diarrhea in, resulting in the development of antibiotic-resistant bacteria, a serious concern for both human and veterinary medicine. In our previous research, we found that *Escherichia coli* was responsible for 85.04% of buffalo calf diarrhea cases, with 69.81% of those isolates being multidrug resistant.²

The commensal *E. coli*, which lives in the intestines of calves, has the ability to acquire resistance genes from other microorganisms or the environment. Thus, it may act as a potential reservoir for the horizontal transmission

of these genes to various bacterial species found in the food chain.³⁻⁵ To control multidrug-resistant *E. coli* in calves, new strategies must be developed in response to the increasing antibiotic resistance. One novel approach to combat antibiotic-resistant *E. coli* is the use of probiotics, which are defined as live microorganisms that, when administered in adequate amounts, provide a health benefit to the host.⁶

Numerous probiotic strains, such as *Bifidobacterium* sp., *Lactobacillus* sp., *Saccharomyces boulardii*, and *Streptococcus thermophilus*, are being used to improve animal health, feed efficiency, weight gain, and immunocompetence.⁷ One of the ideal qualities of an organism to be used as a probiotic species is to adhere and colonize the intestinal lining and must be able to withstand the harsh conditions of bile salt and stomach acid.⁸ The normal gut microbiota is more resilient and stable in the face of population changes. Therefore, choosing species specific probiotic organisms from the gut microbiota may

*Correspondence:

Srivani Moturi. MVSc, PhD

Department of Veterinary Microbiology, N.T.R College of Veterinary Science, Sri Venkateswara Veterinary University, Vijayawada, India

E-mail: srivanimoturi@yahoo.com



This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International (CC BY-NC-SA 4.0) which allows users to read, copy, distribute and make derivative works for non-commercial purposes from the material, as long as the author of the original work is cited properly.

have advantages as autochthonous probiotic species are easy to colonize the intestine, making it easier to maintain the gut microbiota and get rid of enteric infections.⁹

According to previous studies, *Lactobacillus* isolates from fecal samples of dairy cows have been shown to possess both antibacterial properties and the survival qualities required to be potent probiotics.¹⁰ Similar results were found in swine, where probiotics made from autochthonous *Lactobacillus* isolates were more effective.¹¹ Due to the rise in multidrug resistant *E. coli* in buffalo calves, the present research aimed to isolate potential autochthonous *Lactobacillus* species to be used as probiotics against multidrug resistant *E. coli*.

Materials and Methods

Sample collection. From August 2019 to August 2021, a total of 252 rectal swabs were collected from healthy buffalo calves less than 30 days old. These calves were from six districts of Andhra Pradesh. The calves had not received any antibiotics since birth. The swabs were immediately placed into De Man–Rogosa–Sharpe agar (MRS) broth, (HiMedia Laboratories, Kennett Square, USA) containing L-cysteine HCl (0.05%) and Bromocresol green (0.004%) with a low pH of 5.00. They were then incubated at 37.00 °C for 24 to 48 hr in a 5.00% CO₂ incubator.

Isolation and biochemical characterization of *Lactobacillus* species. A loopful of enriched broth culture was streaked on MRS agar with L-cysteine HCl (0.05%) and Bromocresol green (0.004%) plates and incubated at 37.00 °C for 24 to 48 hr in a 5.00% CO₂ incubator. Colonies of dark or light green colour, with a diameter of 2.00 - 3.00 mm and a transparent halo surrounding them were examined for Gram's reaction. Gram-positive rods were then streaked on MRS agar plates and incubated for 48 hr at 37.00 °C in a 5.00% CO₂ incubator. These colonies were further examined for spore staining, motility and biochemical tests such as catalase, indole and nitrate reduction tests.¹² Catalase negative isolates were then inoculated into MRS broth for molecular characterization.

Confirmation of *Lactobacillus* sp. by genus specific polymerase chain reaction (PCR). Whole cell DNA of *Lactobacillus* isolates was extracted using the boiling and snap chilling method¹³ and were subjected to genus-specific PCR using primers Lac 1F (5'-AGCAGTAGGGAATC TTCCA-3') and Lab-0677R, (5'-CACCGCTACACATGGAG-3') targeting the 16S rRNA gene.¹⁴ The reaction mixture consisted of 1.50 µL of DNA template from each isolate, 2.50 µL of 10.00 x Taq buffer (Thermo Fisher Scientific, Waltham, USA), 0.50 µL of 10.00 mM dNTP mix (Thermo Fisher Scientific) 1.50 µL of 25.00 mM MgCl₂ (New England BioLabs, Ipswich, Massachusetts, USA) 1.00 µL of 10.00 pmol µL⁻¹ forward primer, 1.00 µL of 10.00 pmol µL⁻¹ reverse primer, 1.00 µL of 1.00 U µL⁻¹ Taq DNA polymerase (Thermo Fisher Scientific) and 16.00 µL of

nuclease free water to make a total volume of 25.00 µL. The standardized thermal cycling conditions included an initial denaturation at 95.00 °C for 3 min followed by 35 cycles of denaturation at 95.00 °C for 30 sec, annealing at 60.00 °C for 1 min, elongation at 72.00 °C for 1 min and a final elongation at 72.00 °C for 7 min, with an expected amplicon size of 341 bp. DNA from *Lactocaseibacillus rhamnosus*. Microbial Type Culture Collection 1,408 was used as a positive control. The amplified PCR products were then analysed using 1.50% agarose gel electrophoresis and the bands were visualized under an ultraviolet transilluminator (Bio-Rad, Hercules, USA).

Auto-aggregation assay. This test was conducted following the method described Janković *et al.*¹⁵ with slight modifications. *Lactobacillus* cultures grown overnight in MRS broth were harvested by centrifugation at 3,000 rpm for 5 min, then washed and suspended in Phosphate-buffered saline (PBS) to achieve a final optical density of 1.00×10^9 colony-forming unit (CFU) mL⁻¹ at 600 nm. The optical density values were measured with Multiskan GO (Thermo Fisher Scientific) at 5 and 24 hr. Auto-aggregation was calculated as follows:

$$\text{Auto-aggregation (\%)} = [1 - (A_t/A_0)] \times 100$$

where, A_0 represents the absorbance at 0 hr and A_t is the absorbance at 5 and 24 hr.

Cell surface hydrophobicity. *Lactobacillus* cultures grown overnight were centrifuged at 7,500 rpm for 5 min at 4.00 °C, washed twice with PBS buffer and then suspended in the same buffer. The initial absorbance (A_0) at 600 nm was measured. The 2.00 mL of bacterial suspension was then transferred into a round bottom test tube and 0.40 mL of hydrocarbon (N-Hexadecane and xylene; Sigma-Aldrich, Burlington, USA) was added. The tubes were vortexed for 2 min and then left undisturbed for 1 hr to allow the phase separation. The aqueous phase was separated and the absorbance (A_1) was measured at 600 nm. Hydrophobicity was calculated as the percentage decrease ($H\%$) in the absorbance of the bacterial suspension using the following formula:¹⁶

$$H (\%) = [1 - (A_1/A_0)] \times 100$$

Detection of antibacterial activity. The antibacterial activity against multidrug resistant *E. coli* was determined using agar well diffusion method.¹⁷ The degree of inhibition of the tested pathogen was interpreted as high (> 15.00 mm diameter of the zone of inhibition; +++), medium (10.00 - 15.00 mm diameter of zone of inhibition; ++), low (< 10.00 mm diameter of the zone of inhibition; +) and absent (-).¹⁷

Acidic pH tolerance test. Overnight grown *Lactobacillus* cultures were inoculated (1.00 % v/v) into MRS broth that had been previously adjusted to pH values 2.50, and 7.00 (control) with 1.00 N HCl or NaOH (Sisco Research Laboratories Pvt. Ltd., Mumbai, India) incubated at 37.00 °C for 3 hr in a 5.00% CO₂ incubator. The biomass

(CFU mL⁻¹) of each culture obtained in the assays, which were conducted in triplicate, was enumerated on MRS agar incubated anaerobically at 37.00 °C for 24 hr. The reduction in cell count compared with the control tube was assessed as a Log reduction in CFU¹⁸ with slight modifications.

Bile salt tolerance. Overnight grown cultures (1.00 % v/v) were inoculated into MRS broth containing 0.50% (w/v) bovine bile (Sigma-Aldrich) and incubated aerobically at 37.00 °C for 2, 4 and 6 hr.¹⁹ The pH of both control and test cultures was adjusted to 6.00 with 1.00 N HCl or NaOH. The turbidity of the cultures was monitored spectroscopically at 2 hr intervals for growth at 600 nm. The control (blank) consisted of MRS broth without bile.

Antibiotic susceptibility test. Antibiotic susceptibility testing was conducted on selected isolates.²⁰ The bacterial suspension density was adjusted until visible turbidity matched 0.50 McFarland standard (1.50 × 10⁸ CFU mL⁻¹). The inoculum was spread across MRS agar plates. Antibiotic paper discs (HiMedia Laboratories, Mumbai, India) containing ampicillin (10.00 µg), gentamicin (10.00 µg), cefotaxime clavulanic acid (30.00 µg), vancomycin (30.00 µg), tetracycline (30.00 µg), chloramphenicol (30.00 µg), nalidixic acid (30.00 µg), cotrimoxazole (25.00 µg), erythromycin (15.00 µg), penicillin (10.00 U), ciprofloxacin (5.00 µg), clindamycin (2.00 µg), streptomycin (300 µg) and nitrofurantoin (300 µg) were placed on the plates and incubated at 37.00 °C in a 5.00% CO₂ incubator for 24 hr. The diameters of inhibition zone were measured and the results were interpreted as resistant, intermediate, or susceptible according to established standards.²¹

Sequencing and phylogenetic analysis. The 16S rRNA PCR amplified products of different *Lactobacillus* isolates were sequenced using Sanger on a 3,500 genetic analyser (Applied Biosystems, Foster City, USA). To identify the species of the isolate, a similarity search was performed using BLAST in the NCBI database. The phylogenetic analysis of sequences was conducted by MEGA Software (version XI; BioDesign Institute, Tempe, USA).²² A model test was performed to identify the best model that explains the sequence evolution.²³ A neighbour joining (NJ) tree with 1,000 boot strap replications was created using the orthologous 16s rRNA sequence of *B. subtilis* as an outgroup.

Hemolytic activity. The hemolytic activity of isolates was determined using Columbia agar containing 5.00% (w/v) sheep blood (HiMedia Laboratories, Mumbai, India). The plates were then incubated at 37.00 °C for 48 hr. The hemolytic activity of the isolated strains was evaluated and classified based on lysis of red blood cells in the medium surrounding the colonies. Green zones around colonies indicated α-hemolysis, clear zones indicated β-hemolysis, and no zones indicated γ-hemolysis on Columbia blood agar plates. Only strains with γ-hemolysis are considered safe.²⁴

Gelatine liquefaction test. A gelatine medium (Sigma-Aldrich) containing 12.00% gelatine was inoculated with *Lactobacillus* strains at a concentration of 1.00 × 10⁹ CFU mL⁻¹ and incubated for 48 hr at 37.00 °C. Gelatine liquefaction of strains was assessed by storing the medium in a refrigerator for 24 hr and checking whether the gelatine was hydrolysed or not.²⁵

Results

A total of 190 *Lactobacillus* spp. was isolated from 252 samples based on colony morphology, Gram staining (+), spore formation (non-spore formers), motility (non-motile) and different biochemical tests like catalase (-), oxidase (-), indole (-), and nitrate reduction (-) tests. These isolates were further confirmed to be *Lactobacillus* sp. using the genus-specific 16S rRNA PCR (Fig. 1). This study observed a high (100%) isolation rate of *Lactobacillus* sp. from fecal samples of 1 - 2-week age group calves, followed by 97.95% in less than one week age group. In the 2 - 3 weeks and 3 - 4 weeks age groups, 51.78 and 55.81% isolation rates were observed, respectively. The suitability of these isolates for use as probiotic species was further investigated. All of the isolates were tested for auto aggregation and the hydrophobicity of the cell surface using microbial adhesion to hydrocarbons. Among 190 isolates, 57 showed both a high degree of auto-aggregation (> 80.00%) and hydrophobicity (> 60.00%) and were initially selected. The details of the isolates are presented in Table 1. Among the tested hydrocarbons, the maximum adhesion score was observed for N-hexadecane (56.65%) compared to xylene (54.25%).



Fig. 1. Molecular detection of *Lactobacillus* sp. by genus-specific polymerase chain reaction targeting 16s rRNA gene (341bp). Lane 1: MTCC1408, Lanes 2-5: *Lactobacillus* isolates, Lane 6: Negative control, and Lane M: Marker.

Table 1. Auto-aggregation and hydrophobicity (%) of *Lactobacillus* isolates.

Isolate No.	Auto-aggregation	Hydrophobicity		Isolate No.	Auto-aggregation	Hydrophobicity	
		Xylene	N-Hexadecane			Xylene	N-Hexadecane
L10	90.18 ± 0.01	88.52 ± 0.08	89.14 ± 0.05	L103	86.71 ± 0.02	64.37 ± 0.03	75.45 ± 0.09
L12	92.25 ± 0.08	75.80 ± 0.03	80.35 ± 0.01	L108	85.01 ± 0.04	65.73 ± 0.05	75.65 ± 0.03
L15	90.95 ± 0.01	60.30 ± 0.02	61.49 ± 0.03	L109	84.18 ± 0.02	73.37 ± 0.04	76.82 ± 0.04
L21	93.14 ± 0.05	85.63 ± 0.03	75.89 ± 0.05	L125	80.91 ± 0.03	71.13 ± 0.05	87.69 ± 0.05
L33	80.34 ± 0.04	95.49 ± 0.04	62.22 ± 0.05	L126	82.24 ± 0.06	75.70 ± 0.06	78.12 ± 0.07
L34	90.14 ± 0.01	73.34 ± 0.02	87.65 ± 0.03	L134	81.70 ± 0.03	73.72 ± 0.04	77.13 ± 0.06
L35	90.73 ± 0.03	94.22 ± 0.03	84.59 ± 0.05	L135	85.74 ± 0.04	78.69 ± 0.03	69.90 ± 0.02
L37	95.27 ± 0.04	86.62 ± 0.04	89.43 ± 0.02	L136	87.32 ± 0.05	85.31 ± 0.05	88.65 ± 0.03
L38	84.14 ± 0.06	84.56 ± 0.04	82.34 ± 0.04	L143	80.64 ± 0.03	78.21 ± 0.03	84.26 ± 0.04
L39	94.89 ± 0.04	78.28 ± 0.02	86.77 ± 0.05	L144	81.65 ± 0.06	85.34 ± 0.04	88.78 ± 0.05
L42	87.16 ± 0.05	89.76 ± 0.01	92.56 ± 0.09	L145	86.20 ± 0.04	63.25 ± 0.09	86.01 ± 0.07
L43	93.93 ± 0.02	94.46 ± 0.02	95.03 ± 0.02	L162	83.28 ± 0.03	62.82 ± 0.06	63.83 ± 0.04
L44	80.47 ± 0.04	66.09 ± 0.03	93.77 ± 0.05	L163	84.38 ± 0.04	60.31 ± 0.03	72.67 ± 0.05
L45	89.08 ± 0.01	66.42 ± 0.03	64.53 ± 0.05	L169	82.05 ± 0.02	60.70 ± 0.07	61.97 ± 0.06
L51	80.36 ± 0.05	93.11 ± 0.04	94.91 ± 0.03	L171	85.85 ± 0.03	67.34 ± 0.03	69.87 ± 0.04
L57	93.59 ± 0.03	74.81 ± 0.05	88.91 ± 0.03	L173	89.27 ± 0.04	66.05 ± 0.05	70.86 ± 0.08
L58	87.92 ± 0.02	75.43 ± 0.01	94.35 ± 0.04	L177	84.97 ± 0.05	59.61 ± 0.02	70.78 ± 0.03
L59	95.27 ± 0.04	63.12 ± 0.01	62.04 ± 0.04	L178	89.76 ± 0.02	60.12 ± 0.04	68.04 ± 0.04
L60	94.20 ± 0.01	78.37 ± 0.03	65.10 ± 0.03	L182	87.26 ± 0.03	61.40 ± 0.03	69.06 ± 0.06
L63	92.73 ± 0.03	85.67 ± 0.09	89.71 ± 0.04	L187	92.35 ± 0.02	95.27 ± 0.05	95.66 ± 0.05
L66	93.45 ± 0.06	74.25 ± 0.02	68.39 ± 0.04	L191	85.85 ± 0.01	77.51 ± 0.04	75.08 ± 0.07
L68	90.22 ± 0.05	60.63 ± 0.03	73.01 ± 0.05	L196	89.84 ± 0.02	69.03 ± 0.05	62.21 ± 0.06
L70	84.25 ± 0.01	72.39 ± 0.04	68.71 ± 0.03	L202	80.13 ± 0.03	62.41 ± 0.07	68.52 ± 0.05
L71	80.85 ± 0.03	75.74 ± 0.05	73.58 ± 0.01	L203	92.53 ± 0.04	95.68 ± 0.03	95.95 ± 0.07
L73	85.03 ± 0.04	68.00 ± 0.04	72.75 ± 0.03	L204	80.69 ± 0.02	95.50 ± 0.04	95.79 ± 0.05
L80	85.49 ± 0.02	68.44 ± 0.03	60.35 ± 0.04	L207	88.06 ± 0.03	72.40 ± 0.05	65.28 ± 0.07
L94	85.01 ± 0.05	62.85 ± 0.04	64.92 ± 0.03	L209	92.02 ± 0.04	88.25 ± 0.02	74.36 ± 0.06
L99	87.91 ± 0.07	69.15 ± 0.09	85.09 ± 0.06	L211	89.27 ± 0.03	60.56 ± 0.04	62.67 ± 0.05
L100	90.35 ± 0.00	60.80 ± 0.01	60.93 ± 0.05				

The results of the agar well diffusion assay of 57 *Lactobacillus* isolates against multidrug-resistant *E. coli* is presented in Table 2. Of these isolates, 51 exhibited a zone diameter of inhibition (ZDI) greater than >15mm. The isolates L12, L21, L33, L35, L39 and L.143 showed a ZDI less than 15.0 mm.

For probiotic bacteria to survive in the gut, they must be resistant to stomach acidity and have a high tolerance for bile salts. Among the 51 isolates tested, 36 were found to be both acid and bile tolerant. These selected isolates showed a less than a 2-log reduction in cell viability (CFU mL⁻¹) after being incubated at a pH of 2.50 for 3 hr (acid tolerant) and were also tolerant to 0.50% oxgall (bile tolerant). Since the isolates were able to survive in the simulated gastric acid and intestinal environments, we have selected these *Lactobacillus* sp. for further assessment of their antimicrobial activity.

The antibiotic sensitivity test of *Lactobacillus* sp. revealed varying levels of sensitivity to tested antibiotics ranging from 94.44% for ampicillin to 2.78% for vancomycin and streptomycin. All isolates showed 94.44% resistance to vancomycin.

Nucleotide sequencing was performed on 36 *Lactobacillus* isolates which were resistant to bile and acid as well as had > 15.00 mm ZDI on agar well diffusion assay. Among these 22 isolates showed nucleotide sequence

similarity of 99.00% as *Limosilactobacillus fermentum*, six isolates as *Limosilactobacillus reuteri*, four isolates as *Ligilactobacillus agilis*, two isolates as *Lactiplantibacillus plantarum* and one isolate each as *Ligilactobacillus salivarius* and *Limosilactobacillus mucosae*. The hemolysis test revealed that of these 36 isolates five (*L. reuteri* 178, *L. reuteri* 209, *L. fermentum* 182, *L. fermentum* 211, and *L. plantarum* 34) did not exhibit hemolysis (γ-hemolysis) and 31 isolates showed a clear zone of hemolysis (β-hemolysis) around the colonies. Gelatin hydrolysis was not detected in any of the 36 isolates.

The sequences were submitted to the NCBI database with the following accession numbers: *L. fermentum* (OQ283736- OQ283765), *L. plantarum* sub sp. *plantarum* (OQ283773- OQ283774); *L. reuteri* (OQ283780 - OQ283785), *L. mucosae* (OQ283804), *L. agilis* (OQ283800 - OQ283803) and *L. salivarius* (Q283979). The model test revealed that the K2P + G model best explained the sequence evolution. The NJ tree revealed two major clusters (Fig. 2). The first cluster consisted of *L. fermentum*, *L. reuteri*, *L. mucosae* and *L. plantarum* species; with three sub-clusters: *L. fermentum* subgroup, *L. plantarum* subgroup and *L. reuteri* subgroup which included *L. mucosae*. The second cluster included *L. agilis* and *L. salivarius*. All the *Lactobacillus* isolates were grouped distantly from *Bacillus subtilis*.


Table 2. Agar well diffusion test (inhibitory zone) of *Lactobacillus* isolates against multidrug resistant *Escherichia coli*.

Isolate No.	Diameter (mm)	Isolate No.	Diameter (mm)
L10	16.00 ± 0.04	L103	16.00 ± 0.04
L12	14.00 ± 0.01	L108	16.00 ± 0.03
L15	19.00 ± 0.07	L109	19.00 ± 0.03
L21	13.00 ± 0.03	L125	20.00 ± 0.01
L33	14.00 ± 0.09	L126	18.00 ± 0.04
L34	18.00 ± 0.06	L134	18.00 ± 0.05
L35	14.00 ± 0.03	L135	18.00 ± 0.09
L37	18.00 ± 0.01	L136	18.00 ± 0.08
L38	14.00 ± 0.04	L143	14.00 ± 0.01
L39	13.00 ± 0.05	L144	17.00 ± 0.00
L42	22.00 ± 0.01	L145	19.00 ± 0.03
L43	15.00 ± 0.03	L162	18.00 ± 0.03
L44	19.00 ± 0.05	L163	18.00 ± 0.00
L45	20.00 ± 0.06	L169	17.00 ± 0.01
L51	19.00 ± 0.04	L171	15.00 ± 0.02
L57	20.00 ± 0.07	L173	19.00 ± 0.02
L58	16.00 ± 0.09	L177	18.00 ± 0.04
L59	17.00 ± 0.08	L178	17.00 ± 0.06
L60	15.00 ± 0.02	L182	19.00 ± 0.05
L63	23.00 ± 0.04	L187	16.00 ± 0.03
L66	18.00 ± 0.03	L191	20.00 ± 0.06
L68	20.00 ± 0.02	L196	20.00 ± 0.09
L70	18.00 ± 0.03	L202	16.00 ± 0.05
L71	18.00 ± 0.04	L203	15.00 ± 0.05
L73	16.00 ± 0.02	L204	20.00 ± 0.07
L80	17.00 ± 0.02	L207	20.00 ± 0.05
L94	16.00 ± 0.03	L209	20.00 ± 0.02
L99	19.00 ± 0.02	L211	19.00 ± 0.01
L100	17.00 ± 0.01		

Discussion

A total of 190 *Lactobacillus* sp. was isolated and confirmed as *Lactobacillus* based on cultural, morphological, biochemical and molecular testes. Consistent with present findings, several authors earlier isolated and identified *Lactobacillus* as Gram-positive, catalase-negative, non-sporulated rods from fecal samples of different animals.²⁶⁻²⁸ The isolates were further confirmed to be *Lactobacillus* using genus-specific PCR. In neonatal calves, switching their diet from colostrum to whole milk increased the number of milk-using bacteria such as *Lactobacillus*, *Parabacteroides*, and *Bacteroides* in the calf gut²⁹ up to 2 weeks of age. After two to three weeks, calves are gradually introduced to concentrate and hay in addition to milk, which promotes the growth of amylolytic and fibrinolytic bacteria like *Succino-vibrionaceae*, *Fibrobacteraceae*, and *Prevotellaceae* in the developing rumen.³⁰ This may explain the decrease in *Lactobacilli* isolation rates in older calves.

One of the key characteristics of a potent probiotic organism is the adherence to cell surfaces and competitive exclusion of pathogenic bacteria. An increase in hydrophobicity of the cell surface causes enhanced adhesion, and vice versa.³¹ Among the two tested hydro-

**Fig. 2.** Phylogenetic analysis of *Lactobacillus* species.

carbons, N-hexadecane (56.65%) showed the maximum score of adhesion compared to xylene (54.25%). Similar results showing higher adhesion for hexadecane compared to xylene have been reported.³² This study observed that the *Lactobacillus* isolates from calves fed only milk had better auto aggregation than those from calves fed both milk and roughage. Furthermore, isolates from organized dairy farms demonstrated a high ability for auto aggregation compared to isolates from calves kept by individual farmers. The adhesion process is influenced by a number of factors, including surface exopolysaccharides, S-layer protein, and lipoteichoic acid.³³

Out of 57 isolates, 51 exhibited more than 15.00 mm ZDI which is categorised as highly active.³⁴ The organic acids produced by *Lactobacillus* strains reduced the pH of the culture media³⁵ potentially inhibiting the growth of *E. coli*. These organic acids may also act as permeabilizers of outer membrane of Gram-negative bacteria³⁶ enhancing the activity of other antimicrobial metabolites against *E. coli*. Earlier *in vitro* studies have also demonstrated the antimicrobial activity of *Lactobacillus* strains against *E. coli*³⁷ and *Staphylococcus aureus*.³⁸

A potent probiotic bacteria must overcome the physiological barriers of the host, including the hostile acidic environments⁶ and bile toxicity present in the

stomach and duodenum, respectively, in order for the chosen strain to survive in sufficient numbers to express their health-promoting functions in the gut to the best of their ability. The gastric pH of neonatal calves can range from 2.50 to 3.00.³⁹ Therefore, we tested the strains with more than 15.00 mm ZDI for tolerance to acid (pH 2.50) and bile (0.50%). However, considerable variation was observed in acid and bile tolerance among the *Lactobacillus* isolates. Similar findings were also observed in *Lactobacillus* isolates from buffalo calves.²⁶

The antimicrobial susceptibility test revealed that all isolates except *L. fermentum* 66, showed 94.44% resistance to vancomycin. The resistance to vancomycin was previously documented as intrinsic or natural.⁴⁰ According to European Food Safety Authority 2012 guidelines, *Lactobacillus* sp. to be used as a feed additive must be susceptible to ampicillin, streptomycin, gentamicin, erythromycin, clindamycin, tetracycline and chloramphenicol.⁴¹ The observed antimicrobial resistance of *Lactobacillus* sp. against vancomycin, streptomycin and gentamicin in this study may be chromosomally encoded and is an intrinsic feature of *Lactobacillus*, which is non-transferable.^{42,43}

The nucleotide sequencing results of this study revealed that *L. fermentum* is highly prevalent in buffalo calves in this geographic area. The probiotic qualities of *L. fermentum* were discovered in an animal model experiment, including adhesion properties, lifespan extension, immune system strengthening, and other health-promoting capabilities.⁴⁴ Further, it has been found to have synbiotic benefits by increasing antioxidant and immune function in aged mice.⁴⁵ This is consistent with previous findings demonstrating the probiotic qualities of *L. reuteri*⁴⁶ and *L. plantarum*.⁴⁷

Prior to being used as probiotic species *in vivo*, they need to pass tests for other biosafety needs such as hemolysis and gelatine hydrolysis in addition to antibiotic resistance. Hemolysis on blood agar medium occurs when a test bacterium destroys red blood cells by producing α hemolysin, ⁴⁸ resulting in a red transparent liquid. Since hemolysis is associated with pathogenicity, the test probiotic species should be non-hemolytic. In the present study we observed that among 36 *Lactobacillus* isolates *L. reuteri* 178, *L. reuteri* 209, *L. fermentum* 182, *L. fermentum* 211 and *L. plantarum* 34 were non-hemolytic (γ -hemolysis) and none of the 36 isolates showed gelatine hydrolysis. Therefore, these isolates were selected as probiotic species.

In conclusion, based on the results of this study, autochthonous probiotics may be used as feed additives to control multidrug resistant *E. coli* in buffalo calves. Further research is required to determine the therapeutic potential of these probiotics in treating cases of multidrug resistant *E. coli* associated diarrhea in buffalo calves.

Acknowledgments

We are extremely grateful to the Department of Science and Technology - Science and Engineering Research Board, Government of India, for providing the funding necessary to complete this research project. We are also grateful to Sri Venkateswara Veterinary University, Tirupati, for providing the facilities necessary to complete the research project at the Department of Veterinary Microbiology, N.T.R College of Veterinary Science, Gannavaram, India.

Conflict of interest

The authors declare that they have no conflict of interest.

References

1. Axelsson, L. Lactic acid bacteria: classification and physiology. In: Salminen S, Wright AV, Ouwehand A (Eds). Lactic acid bacteria: microbiological and functional aspects. 3rd ed. New York, USA: Marcel Dekker 2004; 1-67.
2. Srivani M, Reddy YN, Subramanyam KV, et al. Prevalence and antimicrobial resistance pattern of Shiga toxinogenic *Escherichia coli* in diarrheic buffalo calves, *Vet World* 2017; 10(7): 774-778.
3. Trobos M, Lester CH, Olsen JE, et al. Natural transfer of sulphonamide and ampicillin resistance between *Escherichia coli* residing in the human intestine. *J Antimicrob. Chemother* 2009; 63(1): 80-86.
4. Ahmed AM, Younis EE, Osman SA, et al. Genetic analysis of antimicrobial resistance in *Escherichia coli* isolated from diarrheic neonatal calves. *Vet Microbiol* 2009; 136(3-4): 397-402.
5. Marshall BM, Levy SB. Food animals and antimicrobials: impacts on human health. *Clin Microbiol Rev* 2011; 24(4): 718-733.
6. World Health Organization. Report of a joint FAO/WHO working group on drafting guidelines for the evaluation of probiotics in food. Available at: <https://www.mhlw.go.jp/file/05-Shingikai-11121000-Iyaku-shokuhinkyoku-Soumuka/0000197343.pdf>. Accessed May 05, 2024.
7. Al-Saiady MY. Effect of probiotic bacteria on immunoglobulin G concentration and other blood components of newborn calves. *J Anim Vet Adv* 2010; 9(3): 604-609.
8. Mahasneh AM, Abbas MM. Probiotics: the possible alternative to disease chemotherapy. In: Darvishi Harzevili F (Ed). *Microbial biotechnology: progress and trends*. 1st ed. Florida, USA: CRC Press 2014; 213-238.
9. Soto LP, Frizzo LS, Bertozzi E, et al. Molecular microbial analysis of *Lactobacillus* strains isolated from the gut of

- calves for potential probiotic use. *Vet Med Int* 2010; 2010: 274987. doi: 10.4061/2010/274987.
10. Adeniyi BA, Adetoye A, Ayeni FA. Antibacterial activities of lactic acid bacteria isolated from cow faeces against potential enteric pathogens. *Afr Health Sci* 2015; 15(3): 888-895.
 11. Balasingham K, Valli C, Radhakrishnan L, et al. Probiotic characterization of lactic acid bacteria isolated from swine intestine. *Vet World* 2017; 10(7): 825-829.
 12. Boone DR, Castenholz RW. *Bergey's manual of systematic bacteriology*. 2nd ed. New York, USA; Springer 2001; 33-38.
 13. Arora S, Agarwal RK, Bist B. Comparison of ELISA and PCR vis-à-vis cultural methods for detecting *Aeromonas* spp. in foods of animal origin. *Int J Food Microbiol* 2006; 106(2): 177-183.
 14. Garcia-Mazcorro JF, Lanerie DJ, Dowd SE, et al. Effect of a multi-species synbiotic formulation on fecal bacterial microbiota of healthy cats and dogs as evaluated by pyrosequencing. *FEMS Microbiol Ecol* 2011; 78(3): 542-554.
 15. Janković T, Frece J, Abram M, et al. Aggregation ability of potential probiotic *Lactobacillus plantarum* strains. *Sanitarno Inženirstvo (IJSER)* 2012; 6(1): 19-24.
 16. Vineetha PG, Tomar S, Saxena VK, et al. Screening of *Lactobacillus* isolates from gastrointestinal tract of guinea fowl for probiotic qualities using in-vitro tests to select species-specific probiotic candidates. *Br Poult Sci* 2016; 57(4): 474-482.
 17. Nair PS. Studies on lactic acid bacteria from tropical fish and shell fish. PhD Thesis. Cochin University of Science and Technology, Cochin, India: 2000.
 18. Guo Z, Wang J, Yan L, et al. *In vitro* comparison of probiotic properties of *Lactobacillus casei* Zhang, a potential new probiotic, with selected probiotic strains. *LWT* 2009; 42(10): 1640-1646.
 19. Khalil R, Mahrous H, El-Halafawy K, et al. Evaluation of the probiotic potential of lactic acid bacteria isolated from faeces of breastfed infants in Egypt. *Afr J Biotechnol* 2007; 6(7): 939-949.
 20. Bauer AW, Kirby MM, Sherris JC, et al. Antibiotic susceptibility testing by a standardized single disk method. *Am J Clin Pathol* 1966; 45(4): 493-496.
 21. Charteris WP, Kelly PM, Morelli L, et al. Development and application of an in vitro methodology to determine the transit tolerance of potentially probiotic *Lactobacillus* and *Bifidobacterium* species in the upper human gastrointestinal tract. *J Appl Microbiol* 1998; 84(5): 759-768.
 22. Tamura K, Stecher G, Kumar S, et al. MEGA11: molecular evolutionary genetics analysis version 11. *Mol Biol Evol* 2021; 38(7): 3022-3027.
 23. Posada D, Crandall KA. MODELTEST: testing the model of DNA substitution. *Bioinformatics* 1998; 14(9): 817-818.
 24. Mangia NP, Saliba L, Deiana P. Functional and safety characterization of autochthonous *Lactobacillus paracasei* FS103 isolated from sheep cheese and its survival in sheep and cow fermented milks during cold storage. *Ann Microbiol* 2019; 69: 161-170.
 25. dela Cruz TEE, Torres JMO. *Gelatin hydrolysis test protocol*. American Society for Microbiology. Washington DC, USA: ASM Press 2012; 1-10.
 26. Singh A, Kumar S, Vinay VV, et al. Autochthonous *Lactobacillus* spp. isolated from Murrah buffalo calves show potential application as probiotic. *Curr Res Biotechnol* 2021; 3: 109-119.
 27. Kumar S, Pattanaik AK, Sharma S, et al. Probiotic potential of a *Lactobacillus* Bacterium of canine faecal origin and its impact on select gut health indices and immune response of dogs. *Probiotics Antimicro Proteins* 2017; 9(3): 262-277.
 28. Tyagi A, Kumar S, Choudhury PK, et al. Conjugated linoleic acid producing potential of lactobacilli isolated from goat (AXB) rumen fluid samples. *Asian-Australas J Anim Sci* 2020; 33(8):1233-1241.
 29. Dias J, Marcondes MI, Noronha MF, et al. Effect of pre-weaning diet on the ruminal archaeal, bacterial, and fungal communities of dairy calves. *Front Microbiol* 2017; 8: 1553. doi: 10.3389/fmicb.2017.01553.
 30. Guzman CE, Bereza-Malcolm LT, De Groef B, et al. Uptake of milk with and without solid feed during the monogastric phase: effect on fibrolytic methanogenic microorganisms in the gastrointestinal tract of calves. *Anim Sci J* 2016; 87(3): 378-388.
 31. Sidira M, Kourkoutas Y, Kanellaki M, et al. *In vitro* study on the cell adhesion ability of immobilized lactobacilli on natural supports. *Food Res Int* 2015; 76(pt 3): 532-539.
 32. Farid W, Masud T, Sohail A, et al. Gastrointestinal transit tolerance, cell surface hydrophobicity, and functional attributes of *Lactobacillus Acidophilus* strains isolated from Indigenous Dahi. *Food Sci Nutr* 2021; 9(9): 5092-5102.
 33. Pan M, Kumaree KK Shah NP. Physiological changes of surface membrane in *Lactobacillus* with prebiotics. *J Food Sci* 2017; 82(3): 744-750.
 34. Chen CC, Lai CC, Huang HL, et al. Antimicrobial activity of *Lactobacillus* species against carbapenem-resistant *Enterobacteriaceae*. *Front Microbiol* 2019; 10: 789. doi: 10.3389/fmicb.2019.00789.
 35. Zhang B, Wang Y, Tan Z, et al. Screening of probiotic activities of *Lactobacilli* strains isolated from traditional Tibetan Qula, a raw yak milk cheese. *Asian-Australas J Anim Sci* 2016; 29(10): 1490-1499.
 36. Alakomi HL, Skyttä E, Saarela M, et al. Lactic acid permeabilizes gram-negative bacteria by disrupting the outer membrane. *Appl Environ Microbiol* 2000; 66(5): 2001-2005.

37. Kumar M, Dhaka P, Vijay D, et al. Antimicrobial effects of *Lactobacillus plantarum* and *Lactobacillus acidophilus* against multidrug-resistant enteroaggregative *Escherichia coli*. *Int J Antimicrob Agents* 2016; 48(3): 265-270.
38. Kang MS, Lim HS, Oh JS, et al. Antimicrobial activity of *Lactobacillus salivarius* and *Lactobacillus fermentum* against *Staphylococcus aureus*. *Pathog Dis* 2017; 75(2). doi: 10.1093/femspd/ftx009.
39. Chandran A, Duary RK, Grover S, et al. Relative expression of bacterial and host specific genes associated with probiotic survival and viability in the mice gut fed with *Lactobacillus plantarum* Lp91. *Microbiol Res* 2013; 168(9): 555-562.
40. Klare I, Konstabel C, Werner G, et al. Antimicrobial susceptibilities of *Lactobacillus*, *Pediococcus* and *Lactococcus* human isolates and cultures intended for probiotic or nutritional use. *J Antimicrob Chemother* 2007; 59(5): 900-912.
41. Marchwińska K, Gwiazdowska D. Isolation and probiotic potential of lactic acid bacteria from swine feces for feed additive composition. *Arch Microbiol* 2021; 204(1): 61. doi: 10.1007/s00203-021-02700-0.
42. Tulini FL, Winkelströter LK, De Martinis EC. Identification and evaluation of the probiotic potential of *Lactobacillus paraplantarum* FT259, a bacteriocinogenic strain isolated from Brazilian semi-hard artisanal cheese. *Anaerobe* 2013; 22: 57-63.
43. Casarotti SN, Carneiro BM, Todorov SD, et al. *In vitro* assessment of safety and probiotic potential characteristics of *Lactobacillus* strains isolated from water buffalo mozzarella cheese. *Ann Microbiol* 2017; 67: 289-301.
44. Park MR, Ryu S, Maburutse BE, et al. Probiotic. *Lactobacillus fermentum* strain JDFM216 stimulates the longevity and immune response of *Caenorhabditis elegans* through a nuclear hormone receptor. *Sci Rep* 2018; 8(1): 7441. doi: 10.1038/s41598-018-25333-8.
45. Sharma R, Kumari M, Kumari A, et al. Diet supplemented with phytochemical epigallocatechin gallate and probiotic *Lactobacillus fermentum* confers second generation synbiotic effects by modulating cellular immune responses and antioxidant capacity in aging mice. *Eur J Nutr* 2019; 58(7): 2943-2957.
46. Mu Q, Tavella VJ, Luo XM. Role of *Lactobacillus reuteri* in human health and diseases. *Front Microbiol* 2018; 9: 757. doi: 10.3389/fmicb.2018.00757.
47. Fidanza M, Panigrahi P, Kollmann TR. *Lactiplantibacillus plantarum*-nomad and ideal probiotic. *Front Microbiol* 2021; 12: 712236. doi: 10.3389/fmicb.2021.712236.
48. Gottschalk MG, Lacouture S, Dubreuil JD. Characterization of *Streptococcus suis* capsular type 2 haemolysin. *Microbiology (Reading)* 1995; 141(Pt 1): 189-195.